

Publications in which pimonidazole HCl was used as a hypoxia marker.

J. A Raleigh, PhD
Hypoxyprobe, Inc
Updated August 2011

2011

1. Boulton JK, Walker-Samuel S, Jamin Y, Leiper JM, Whitley GS, Robinson SP. Active site mutant dimethylarginine dimethylaminohydrolase 1 expression confers an intermediate tumour phenotype in C6 gliomas. *J Pathol* 2011.
2. Busk M, Walenta S, Mueller-Klieser W, et al. Inhibition of tumor lactate oxidation: Consequences for the tumor microenvironment. *Radiother Oncol* 2011; 99: 404-11.
3. Chaudhuri S, McCullough SS, Hennings L, et al. Acetaminophen hepatotoxicity and HIF-1alpha induction in acetaminophen toxicity in mice occurs without hypoxia. *Toxicol Appl Pharmacol* 2011; 252: 211-20.
4. Donaldson SB, Betts G, Bonington SC, et al. Perfusion Estimated with Rapid Dynamic Contrast-Enhanced Magnetic Resonance Imaging Correlates Inversely with Vascular Endothelial Growth Factor Expression and Pimonidazole Staining in Head-and-Neck Cancer: A Pilot Study. *Int J Radiat Oncol Biol Phys* 2011.
5. Egeland TA, Gulliksrud K, Gaustad JV, Mathiesen B, Rofstad EK. Dynamic contrast-enhanced-MRI of tumor hypoxia. *Magn Reson Med* 2011.
6. Erba P, Ogawa R, Ackermann M, et al. Angiogenesis in wounds treated by microdeformational wound therapy. *Ann Surg* 2011; 253: 402-9.
7. Gillies RM, Robinson SP, McPhail LD, Carter ND, Murray JF. Immunohistochemical assessment of intrinsic and extrinsic markers of hypoxia in reproductive tissue: differential expression of HIF1alpha and HIF2alpha in rat oviduct and endometrium. *J Mol Histol* 2011; 42: 341-54.
8. Grahl N, Puttikamonkul S, Macdonald JM, et al. In vivo Hypoxia and a Fungal Alcohol Dehydrogenase Influence the Pathogenesis of Invasive Pulmonary Aspergillosis. *PLoS Pathog* 2011; 7: e1002145.
9. Gulliksrud K, Hompland T, Galappathi K, Rofstad EK. Assessment of tumor hypoxia and interstitial fluid pressure by gadomelitol-based dynamic contrast-enhanced magnetic resonance imaging. *Radiother Oncol* 2011.
10. Gulliksrud K, Ovrebo KM, Mathiesen B, Rofstad EK. Differentiation between hypoxic and non-hypoxic experimental tumors by dynamic contrast-enhanced magnetic resonance imaging. *Radiother Oncol* 2011; 98: 360-4.
11. Harris NR, Carter PR, Yadav AS, et al. Relationship between inflammation and tissue hypoxia in a mouse model of chronic colitis. *Inflamm Bowel Dis* 2011; 17: 742-6.
12. Heng Y, Seah PG, Siew JY, et al. Mycobacterium tuberculosis infection induces hypoxic lung lesions in the rat. *Tuberculosis (Edinb)* 2011; 91: 339-41.
13. Hennings LJ, Flores R, Roberson PK, et al. Persistent penumbra in a rabbit stroke model: incidence and histologic characteristics. *Stroke Res Treat* 2011; 2011: 764830.

14. Hindryckx P, Devisscher L, Laukens D, Venken K, Peeters H, De Vos M. Intrarectal administration of oxygenated perfluorodecalin promotes healing of murine colitis by targeting inflammatory hypoxia. *Lab Invest* 2011.
15. Kang YJ, Jo JO, Cho MK, Yu HS, Ock MS, Cha HJ. *Trichinella spiralis* infection induces angiogenic factor thymosin beta4 expression. *Vet Parasitol* 2011.
16. Maftai CA, Bayer C, Shi K, Astner ST, Vaupel P. Changes in the fraction of total hypoxia and hypoxia subtypes in human squamous cell carcinomas upon fractionated irradiation: Evaluation using pattern recognition in microcirculatory supply units. *Radiother Oncol* 2011.
17. Maftai CA, Bayer C, Shi K, Astner ST, Vaupel P. Quantitative assessment of hypoxia subtypes in microcirculatory supply units of malignant tumors using (immuno-)fluorescence techniques. *Strahlenther Onkol* 2011; 187: 260-6.
18. Maftai CA, Shi K, Bayer C, Astner ST, Vaupel P. Comparison of (immuno-)fluorescence data with serial [(18)F]Fmiso PET/CT imaging for assessment of chronic and acute hypoxia in head and neck cancers. *Radiother Oncol* 2011; 99: 412-7.
19. Mees G, Vangestel C, Dierckx R, et al. Metabolic correlates of tumour hypoxia in malignant canine mammary carcinoma. *Res Vet Sci* 2011.
20. Nagelkerke A, Mujcic H, Bussink J, et al. Hypoxic regulation and prognostic value of LAMP3 expression in breast cancer. *Cancer* 2011; 117: 3670-81.
21. Nijkamp MM, Hoogsteen IJ, Span PN, et al. Spatial relationship of phosphorylated epidermal growth factor receptor and activated AKT in head and neck squamous cell carcinoma. *Radiother Oncol* 2011.
22. Oehler C, O'Donoghue JA, Russell J, et al. 18F-fluoromisonidazole PET imaging as a biomarker for the response to 5,6-dimethylxanthenone-4-acetic acid in colorectal xenograft tumors. *J Nucl Med* 2011; 52: 437-44.
23. Olsson R, Carlsson PO. A low-oxygenated subpopulation of pancreatic islets constitutes a functional reserve of endocrine cells. *Diabetes* 2011; 60: 2068-75.
24. Olsson R, Olerud J, Pettersson U, Carlsson PO. Increased Numbers of Low-Oxygenated Pancreatic Islets After Intraportal Islet Transplantation. *Diabetes* 2011.
25. Orlova AG, Maslennikova AV, Golubiatnikov G, et al. [Noninvasive estimation of the oxygen state of experimental tumor by diffuse optical spectroscopy]. *Biofizika* 2011; 56: 349-55.
26. Rademakers SE, Lok J, van der Kogel AJ, Bussink J, Kaanders JH. Metabolic markers in relation to hypoxia; staining patterns and colocalization of pimonidazole, HIF-1alpha, CAIX, LDH-5, GLUT-1, MCT1 and MCT4. *BMC Cancer* 2011; 11: 167.
27. Sato Y, Endo H, Okuyama H, et al. Cellular hypoxia of pancreatic beta-cells due to high levels of oxygen consumption for insulin secretion in vitro. *J Biol Chem* 2011; 286: 12524-32.
28. Silvola JM, Saraste A, Forsback S, et al. Detection of hypoxia by [18F]EF5 in atherosclerotic plaques in mice. *Arterioscler Thromb Vasc Biol* 2011; 31: 1011-5.
29. Takiyama Y, Harumi T, Watanabe J, et al. Tubular injury in a rat model of type 2 diabetes is prevented by metformin: a possible role of HIF-1alpha expression and oxygen metabolism. *Diabetes* 2011; 60: 981-92.
30. Ueda M, Kudo T, Mutou Y, et al. Evaluation of [(125) I]IPOS as a molecular imaging probe for hypoxia-inducible factor-1-active regions in a tumor: Comparison among SPECT/CT imaging, autoradiography, and immunohistochemistry. *Cancer Sci* 2011.

31. Vigneswaran N, Wu J, Song A, Annapragada A, Zacharias W. Hypoxia-induced autophagic response is associated with aggressive phenotype and elevated incidence of metastasis in orthotopic immunocompetent murine models of head and neck squamous cell carcinomas (HNSCC). *Exp Mol Pathol* 2011; 90: 215-25.
32. Wright WS, McElhatten RM, Harris NR. Increase in retinal hypoxia-inducible factor-2alpha, but not hypoxia, early in the progression of diabetes in the rat. *Exp Eye Res* 2011.
33. Xing D, Liu L, Marti GP, et al. Hypoxia and hypoxia-inducible factor in the burn wound. *Wound Repair Regen* 2011; 19: 205-13.
34. Yaromina A, Kroeber T, Meinzer A, et al. Exploratory study of the prognostic value of microenvironmental parameters during fractionated irradiation in human squamous cell carcinoma xenografts. *Int J Radiat Oncol Biol Phys* 2011; 80: 1205-13.
35. Yeh KY, Yeh M, Polk P, Glass J. Hypoxia-inducible factor-2alpha and iron absorptive gene expression in Belgrade rat intestine. *Am J Physiol Gastrointest Liver Physiol* 2011; 301: G82-90.
36. Yim HE, Kim JH, Yoo KH, Bae IS, Hong YS, Lee JW. Spironolactone and enalapril differentially up-regulate the expression of VEGF and heme oxygenase-1 in the neonatal rat kidney. *Pediatr Res* 2011; 69: 378-83.
37. Zaleska K, Bruechner K, Baumann M, Zips D, Yaromina A. Tumour-infiltrating CD11b+ myelomonocytes and response to fractionated irradiation of human squamous cell carcinoma (hSCC) xenografts. *Radiother Oncol* 2011.
38. Zhao D, Chang CH, Kim JG, Liu H, Mason RP. In vivo near-infrared spectroscopy and magnetic resonance imaging monitoring of tumor response to combretastatin A-4-phosphate correlated with therapeutic outcome. *Int J Radiat Oncol Biol Phys* 2011; 80: 574-81.

2010

1. Araujo AP, Frezza TF, Allegretti SM, Giorgio S. Hypoxia, hypoxia-inducible factor-1alpha and vascular endothelial growth factor in a murine model of *Schistosoma mansoni* infection. *Exp Mol Pathol* 2010.
2. Brown AP, Chung EJ, Urick ME, et al. Evaluation of the fullerene compound DF-1 as a radiation protector. *Radiat Oncol* 2010; 5: 34.
3. Busk M, Munk OL, Jakobsen S, et al. Assessing hypoxia in animal tumor models based on pharmacokinetic analysis of dynamic FAZA PET. *Acta Oncol* 2010; 49: 922-33.
4. Fokas E, Hanze J, Kamlah F, et al. Irradiation-dependent effects on tumor perfusion and endogenous and exogenous hypoxia markers in an A549 xenograft model. *Int J Radiat Oncol Biol Phys* 2010; 77: 1500-8.
5. Graves EE, Vilalta M, Cecic IK, et al. Hypoxia in Models of Lung Cancer: Implications for Targeted Therapeutics. *Clin Cancer Res* 2010; 16: 4843-52.
6. Harris NR, Carter PR, Yadav AS, et al. Relationship between inflammation and tissue hypoxia in a mouse model of chronic colitis. *Inflamm Bowel Dis* 2010.
7. Herrick AL, Gorodkin R, Jeziorska M, Stratford IJ. Testing for hypoxia in forearm skin of patients with systemic sclerosis, assessed by pimonidazole. *J Rheumatol* 2010; 37: 1968-9.

8. Hindryckx P, Waeytens A, Laukens D, et al. Absence of placental growth factor blocks dextran sodium sulfate-induced colonic mucosal angiogenesis, increases mucosal hypoxia and aggravates acute colonic injury. *Lab Invest* 2010; 90: 566-76.
9. Hoeben BA, Kaanders JH, Franssen GM, et al. PET of hypoxia with 89Zr-labeled cG250-F(ab')₂ in head and neck tumors. *J Nucl Med* 2010; 51: 1076-83.
10. Huang RQ, Cheng HL, Zhao XD, et al. Preliminary study on the effect of trauma-induced secondary cellular hypoxia in brain injury. *Neurosci Lett* 2010; 473: 22-7.
11. Javvadi P, Hertan L, Kosoff R, et al. Thioredoxin reductase-1 mediates curcumin-induced radiosensitization of squamous carcinoma cells. *Cancer Res* 2010; 70: 1941-50.
12. Kimura M, Rabbani Z, Mouraviev V, et al. Role of vitamin D(3) as a sensitizer to cryoablation in a murine prostate cancer model: preliminary in vivo study. *Urology* 2010; 76: 764 e14-20.
13. Kimura M, Rabbani Z, Mouraviev V, et al. Morphology of hypoxia following cryoablation in a prostate cancer murine model: its relationship to necrosis, apoptosis and, microvessel density. *Cryobiology* 2010; 61: 148-54.
14. Kurobe H, Urata M, Ueno M, et al. Role of hypoxia-inducible factor 1alpha in T cells as a negative regulator in development of vascular remodeling. *Arterioscler Thromb Vasc Biol* 2010; 30: 210-7.
15. Laifenfeld D, Gilchrist A, Drubin D, et al. The role of hypoxia in 2-butoxyethanol-induced hemangiosarcoma. *Toxicol Sci* 2010; 113: 254-66.
16. Lee DY-W, Ji X-S, Raleigh JA, inventors; Natural Pharmacia International, Inc, assignee. Hypoxia selective weakly basic 2-nitroimidazole delivery agents and methods of use thereof. USA patent 7,842,278 B2. 2010 Nov 30, 2010.
17. Lee JC, Kinniry PA, Arguiri E, et al. Dietary curcumin increases antioxidant defenses in lung, ameliorates radiation-induced pulmonary fibrosis, and improves survival in mice. *Radiat Res* 2010; 173: 590-601.
18. Lehman S, Koumenis C. The role of autophagy as a mechanism of cytotoxicity by the clinically used agent MDA-7/IL-24. *Cancer Biol Ther* 2010; 9.
19. Leno-Duran E, Hatta K, Bianco J, et al. Fetal-placental hypoxia does not result from failure of spiral arterial modification in mice. *Placenta* 2010; 31: 731-7.
20. Li XF, Ma Y, Sun X, Humm JL, Ling CC, O'Donoghue JA. High 18F-FDG uptake in microscopic peritoneal tumors requires physiologic hypoxia. *J Nucl Med* 2010; 51: 632-8.
21. Li XF, Sun X, Ma Y, et al. Detection of hypoxia in microscopic tumors using 131I-labeled iodo-azomycin galactopyranoside (131I-IAZGP) digital autoradiography. *Eur J Nucl Med Mol Imaging* 2010; 37: 339-48.
22. Maity A, Koumenis C. Location, Location, Location-Makes All the Difference for Hypoxia in Lung Tumors. *Clin Cancer Res* 2010; 16: 4685-7.
23. Maslennikova AV, Orlova AG, Golubiatnikov GY, et al. Comparative study of tumor hypoxia by diffuse optical spectroscopy and immunohistochemistry in two tumor models. *J Biophotonics* 2010.
24. Masunaga S, Nagasawa H, Liu Y, et al. Evaluation of the radiosensitivity of the oxygenated tumor cell fractions in quiescent cell populations within solid tumors. *Radiat Res* 2010; 174: 459-66.
25. Matsuzaki S, Jardon K, Maleysson E, et al. Carbon dioxide pneumoperitoneum, intraperitoneal pressure, and peritoneal tissue hypoxia: a mouse study with controlled respiratory support. *Surg Endosc* 2010.

26. McGee MC, Hamner JB, Williams RF, et al. Improved intratumoral oxygenation through vascular normalization increases glioma sensitivity to ionizing radiation. *Int J Radiat Oncol Biol Phys* 2010; 76: 1537-45.
27. McPhail LD, Robinson SP. Intrinsic susceptibility MR imaging of chemically induced rat mammary tumors: relationship to histologic assessment of hypoxia and fibrosis. *Radiology* 2010; 254: 110-8.
28. Moon JW, Kim YJ, Khwarg SI, Chung H, Yu HG. Chorioretinal ischemia and angiogenic milieu following photodynamic therapy. *Curr Eye Res* 2010; 35: 314-21.
29. Morchel P, Melkus G, Yaromina A, et al. Correlating quantitative MR measurements of standardized tumor lines with histological parameters and tumor control dose. *Radiother Oncol* 2010; 96: 123-30.
30. Morelli A, Filippi S, Comeglio P, et al. Acute vardenafil administration improves bladder oxygenation in spontaneously hypertensive rats. *J Sex Med* 2010; 7: 107-20.
31. Mowat FM, Luhmann UF, Smith AJ, et al. HIF-1alpha and HIF-2alpha are differentially activated in distinct cell populations in retinal ischaemia. *PLoS One* 2010; 5: e11103.
32. Nilsson MF, Danielsson C, Skold AC, et al. Improved methodology for identifying the teratogenic potential in early drug development of hERG channel blocking drugs. *Reprod Toxicol* 2010; 29: 156-63.
33. Olive PL, Banuelos CA, Durand RE, Kim JY, Aquino-Parsons C. Endogenous and radiation-induced expression of gammaH2AX in biopsies from patients treated for carcinoma of the uterine cervix. *Radiother Oncol* 2010; 94: 82-9.
34. Pina Y, Houston SK, Murray T, et al. Focal, Periocular Delivery of 2-deoxy-D-glucose as an Adjuvant to Chemotherapy for the Treatment of Advanced Retinoblastoma. *Invest Ophthalmol Vis Sci* 2010.
35. Prasad P, Li LP, Halter S, Cabray J, Ye M, Battle D. Evaluation of Renal Hypoxia in Diabetic Mice by BOLD MRI. *Invest Radiol* 2010.
36. Rabbani ZN, Mi J, Zhang Y, et al. Hypoxia inducible factor 1alpha signaling in fractionated radiation-induced lung injury: role of oxidative stress and tissue hypoxia. *Radiat Res* 2010; 173: 165-74.
37. Robinson MA, Tuttle SW, Otto CM, Koch CJ. pO(2)-dependent NO production determines OPPC activity in macrophages. *Free Radic Biol Med* 2010; 48: 189-95.
38. Santiago A, Eicheler W, Bussink J, et al. Effect of cetuximab and fractionated irradiation on tumour micro-environment. *Radiother Oncol* 2010.
39. Schmeding M, Rademacher S, Boas-Knoop S, et al. rHuEPO reduces ischemia-reperfusion injury and improves survival after transplantation of fatty livers in rats. *Transplantation* 2010; 89: 161-8.
40. Shynlova O, Dorogin A, Lye SJ. Stretch-induced uterine myocyte differentiation during rat pregnancy: involvement of caspase activation. *Biol Reprod* 2010; 82: 1248-55.
41. Simonsen TG, Gaustad JV, Rofstad EK. Development of hypoxia in a preclinical model of tumor micrometastases. *Int J Radiat Oncol Biol Phys* 2010; 76: 879-88.
42. Wright WS, McElhatten RM, Messina JE, Harris NR. Hypoxia and the expression of HIF-1alpha and HIF-2alpha in the retina of streptozotocin-injected mice and rats. *Exp Eye Res* 2010; 90: 405-12.

43. Yaromina A, Thames H, Zhou X, et al. Radiobiological hypoxia, histological parameters of tumour microenvironment and local tumour control after fractionated irradiation. *Radiother Oncol* 2010; 96: 116-22.
44. Yim HE, Kim JH, Yoo KH, et al. Spironolactone, but not enalapril, potentiates hypoxia-inducible factor-1 alpha and Ets-1 expression in newborn rat kidney. *J Physiol Pharmacol* 2010; 61: 73-81.
45. Young RJ, Moller A. Immunohistochemical detection of tumour hypoxia. *Methods Mol Biol* 2010; 611: 151-9.

2009

1. Ahlskog JK, Schliemann C, Marlind J, et al. Human monoclonal antibodies targeting carbonic anhydrase IX for the molecular imaging of hypoxic regions in solid tumours. *Br J Cancer* 2009; 101: 645-57.
2. Bartoli M, Al-Shabrawey M, Labazi M, et al. HMG-CoA reductase inhibitors (statin) prevents retinal neovascularization in a model of oxygen-induced retinopathy. *Invest Ophthalmol Vis Sci* 2009; 50: 4934-40.
3. Boutrid H, Pina Y, Cebulla C, et al. Vessel Targeting Increases Hypoxia in a Murine Model of Retinoblastoma. *Invest Ophthalmol Vis Sci* 2009.
4. Boutrid H, Pina Y, Cebulla CM, et al. Increased hypoxia following vessel targeting in a murine model of retinoblastoma. *Invest Ophthalmol Vis Sci* 2009; 50: 5537-43.
5. Bruechner K, Bergmann R, Santiago A, et al. Comparison of [(18)F]FDG uptake and distribution with hypoxia and proliferation in FaDu human squamous cell carcinoma (hSCC) xenografts after single dose irradiation. *Int J Radiat Biol* 2009; 1-9.
6. Busk M, Horsman MR, Jakobsen S, et al. Can hypoxia-PET map hypoxic cell density heterogeneity accurately in an animal tumor model at a clinically obtainable image contrast? *Radiother Oncol* 2009; 92: 429-36.
7. Carlin S, Pugachev A, Sun X, et al. In vivo characterization of a reporter gene system for imaging hypoxia-induced gene expression. *Nucl Med Biol* 2009; 36: 821-31.
8. Chen J, Connor KM, Aderman CM, Willett KL, Aspegren OP, Smith LE. Suppression of retinal neovascularization by erythropoietin siRNA in a mouse model of proliferative retinopathy. *Invest Ophthalmol Vis Sci* 2009; 50: 1329-35.
9. Cho H, Ackerstaff E, Carlin S, et al. Noninvasive multimodality imaging of the tumor microenvironment: registered dynamic magnetic resonance imaging and positron emission tomography studies of a preclinical tumor model of tumor hypoxia. *Neoplasia* 2009; 11: 247-59, 2p following 59.
10. Du J, Fan YY, Hitomi H, et al. Mineralocorticoid receptor blockade and calcium channel blockade have different renoprotective effects on glomerular and interstitial injury in rats. *Am J Physiol Renal Physiol* 2009.
11. Ellingsen C, Natvig I, Gaustad JV, Gulliksrud K, Egeland TA, Rofstad EK. Human cervical carcinoma xenograft models for studies of the physiological microenvironment of tumors. *J Cancer Res Clin Oncol* 2009.
12. Fujimoto S, Satoh M, Nagasu H, Horike H, Sasaki T, Kashihara N. Azelnidipine exerts renoprotective effects by improvement of renal microcirculation in angiotensin II infusion rats. *Nephrol Dial Transplant* 2009.

13. Hernandez C, Santamatilde E, McCreath KJ, et al. Induction of trefoil factor (TFF)1, TFF2 and TFF3 by hypoxia is mediated by hypoxia inducible factor-1: implications for gastric mucosal healing. *Br J Pharmacol* 2009; 156: 262-72.
14. Hoogsteen IJ, Lok J, Marres HA, et al. Hypoxia in larynx carcinomas assessed by pimonidazole binding and the value of CA-IX and vascularity as surrogate markers of hypoxia. *Eur J Cancer* 2009; 45: 2906-14.
15. Huang G, Chen L. Discrepancies between antiangiogenic and antitumor effects of recombinant human endostatin. *Cancer Biother Radiopharm* 2009; 24: 589-96.
16. Kao YH, Jawan B, Goto S, et al. Serum factors potentiate hypoxia-induced hepatotoxicity in vitro through increasing transforming growth factor-beta1 activation and release. *Cytokine* 2009; 47: 11-22.
17. Kim HK, Bian H, Aya-ay J, Garces A, Morgan EF, Gilbert SR. Hypoxia and HIF-1alpha expression in the epiphyseal cartilage following ischemic injury to the immature femoral head. *Bone* 2009; 45: 280-8.
18. Kizaka-Kondoh S, Konse-Nagasawa H. Significance of nitroimidazole compounds and hypoxia-inducible factor-1 for imaging tumor hypoxia. *Cancer Sci* 2009; 100: 1366-73.
19. Kudo T, Ueda M, Kuge Y, et al. Imaging of HIF-1-active tumor hypoxia using a protein effectively delivered to and specifically stabilized in HIF-1-active tumor cells. *J Nucl Med* 2009; 50: 942-9.
20. Ley EJ, Scehnet J, Park R, et al. The in vivo effect of propranolol on cerebral perfusion and hypoxia after traumatic brain injury. *J Trauma* 2009; 66: 154-9; discussion 9-61.
21. Lin WY, Mannikarottu A, Li S, et al. Correlation of in vivo bladder blood flow measurements with tissue hypoxia. *World J Urol* 2009.
22. Liu Y, Kudo K, Abe Y, et al. Inhibition of Transforming Growth Factor-beta, Hypoxia-inducible Factor-1alpha and Vascular Endothelial Growth Factor Reduced Late Rectal Injury Induced by Irradiation. *J Radiat Res (Tokyo)* 2009.
23. Ma J, Waxman DJ. Dominant effect of antiangiogenesis in combination therapy involving cyclophosphamide and axitinib. *Clin Cancer Res* 2009; 15: 578-88.
24. Mazzone M, Dettori D, Leite de Oliveira R, et al. Heterozygous deficiency of PHD2 restores tumor oxygenation and inhibits metastasis via endothelial normalization. *Cell* 2009; 136: 839-51.
25. Newbold K, Castellano I, Charles-Edwards E, et al. An exploratory study into the role of dynamic contrast-enhanced magnetic resonance imaging or perfusion computed tomography for detection of intratumoral hypoxia in head-and-neck cancer. *Int J Radiat Oncol Biol Phys* 2009; 74: 29-37.
26. Oh M, Tanaka T, Kobayashi M, et al. Radio-copper-labeled Cu-ATSM: an indicator of quiescent but clonogenic cells under mild hypoxia in a Lewis lung carcinoma model. *Nucl Med Biol* 2009; 36: 419-26.
27. Rabbani ZN, Spasojevic I, Zhang X, et al. Antiangiogenic action of redox-modulating Mn(III) meso-tetrakis(N-ethylpyridinium-2-yl)porphyrin, MnTE-2-PyP(5+), via suppression of oxidative stress in a mouse model of breast tumor. *Free Radic Biol Med* 2009.
28. Rosenberger C, Rosen S, Paliege A, Heyman SN. Pimonidazole adduct immunohistochemistry in the rat kidney: detection of tissue hypoxia. *Methods Mol Biol* 2009; 466: 161-74.

29. Russell J, Carlin S, Burke SA, Wen B, Yang KM, Ling CC. Immunohistochemical detection of changes in tumor hypoxia. *Int J Radiat Oncol Biol Phys* 2009; 73: 1177-86.
30. Schelshorn DW, Schneider A, Kuschinsky W, et al. Expression of hemoglobin in rodent neurons. *J Cereb Blood Flow Metab* 2009; 29: 585-95.
31. Schmeding M, Hunold G, Ariyakhagorn V, et al. Erythropoietin reduces ischemia-reperfusion injury after liver transplantation in rats. *Transpl Int* 2009; 22: 738-46.
32. Shibata R, Ueda S, Yamagishi S, et al. Involvement of asymmetric dimethylarginine (ADMA) in tubulointerstitial ischaemia in the early phase of diabetic nephropathy. *Nephrol Dial Transplant* 2009; 24: 1162-9.
33. Vignozzi L, Filippi S, Morelli A, et al. Cavernous neurotomy in the rat is associated with the onset of an overt condition of hypogonadism. *J Sex Med* 2009; 6: 1270-83.
34. Wen QL, Meng MB, Yang B, et al. Endostar, a recombinant humanized endostatin, enhances the radioresponse for human nasopharyngeal carcinoma and human lung adenocarcinoma xenografts in mice. *Cancer Sci* 2009; 100: 1510-9.
35. Wijffels KI, Hoogsteen IJ, Lok J, et al. No detectable hypoxia in malignant salivary gland tumors: preliminary results. *Int J Radiat Oncol Biol Phys* 2009; 73: 1319-25.
36. Xia CY, Li L, Liu HM, Cong WM. High expression of angiotensin-converting enzyme and angiotensin-converting enzyme 2 in preservation injury after liver transplantation in rats. *Hepato Res* 2009.
37. Yang KM, Russell J, Lupu ME, et al. Atrasentan (ABT-627) enhances perfusion and reduces hypoxia in a human tumor xenograft model. *Cancer Biol Ther* 2009; 8: 1940-6.
38. Yaromina A, Eckardt A, Zips D, et al. Core needle biopsies for determination of the microenvironment in individual tumours for longitudinal radiobiological studies. *Radiother Oncol* 2009.
39. Yaromina A, Quennet V, Zips D, et al. Co-localisation of hypoxia and perfusion markers with parameters of glucose metabolism in human squamous cell carcinoma (hSCC) xenografts. *Int J Radiat Biol* 2009; 85: 972-80.
40. Zaghoul N, Hernandez SL, Bae JO, et al. Vascular endothelial growth factor blockade rapidly elicits alternative proangiogenic pathways in neuroblastoma. *Int J Oncol* 2009; 34: 401-7.

2008

1. Bache M, Kappler M, Said HM, Staab A, Vordermark D. Detection and specific targeting of hypoxic regions within solid tumors: current preclinical and clinical strategies. *Curr Med Chem* 2008; 15: 322-38.
2. Bartoli M, Al-Shabrawey M, Labazi M, et al. HMG-CoA reductase inhibitors (statins) prevents retinal neovascularization in a model of oxygen-induced retinopathy. *Invest Ophthalmol Vis Sci* 2008.
3. Bayer C, Schilling D, Hoetzel J, et al. PAI-1 levels predict response to fractionated irradiation in 10 human squamous cell carcinoma lines of the head and neck. *Radiother Oncol* 2008; 86: 361-8.
4. Bouchentouf M, Benabdallah BF, Bigey P, Yau TM, Scherman D, Tremblay JP. Vascular endothelial growth factor reduced hypoxia-induced death of human myoblasts and improved their engraftment in mouse muscles. *Gene Ther* 2008; 15: 404-14.

5. Boutrid H, Jockovich ME, Murray TG, et al. Targeting hypoxia, a novel treatment for advanced retinoblastoma. *Invest Ophthalmol Vis Sci* 2008; 49: 2799-805.
6. Busk M, Horsman MR, Jakobsen S, et al. Imaging Hypoxia in Xenografted and Murine Tumors With (18)F-Fluoroazomycin Arabinoside: A Comparative Study Involving microPET, Autoradiography, Po(2)-Polarography, and Fluorescence Microscopy. *Int J Radiat Oncol Biol Phys* 2008; 70: 1202-12.
7. Busk M, Horsman MR, Kristjansen PE, van der Kogel AJ, Bussink J, Overgaard J. Aerobic glycolysis in cancers: implications for the usability of oxygen-responsive genes and fluorodeoxyglucose-PET as markers of tissue hypoxia. *Int J Cancer* 2008; 122: 2726-34.
8. Busk M, Horsman MR, Overgaard J. Resolution in PET hypoxia imaging: voxel size matters. *Acta Oncol* 2008; 47: 1201-10.
9. Bussink J, van der Kogel AJ, Kaanders JH. Patterns and levels of hypoxia in head and neck squamous cell carcinomas and their relationship to patient outcome: in regard to Evans et al. (*Int J Radiat Oncol Biol Phys* 2007;69:1024-1031). *Int J Radiat Oncol Biol Phys* 2008; 70: 1616.
10. Bussink J, van der Kogel AJ, Kaanders JH. Activation of the PI3-K/AKT pathway and implications for radioresistance mechanisms in head and neck cancer. *Lancet Oncol* 2008; 9: 288-96.
11. Choi YH, Cowan DB, Nathan M, et al. Myocardial hypertrophy overrides the angiogenic response to hypoxia. *PLoS ONE* 2008; 3: e4042.
12. Dai M, Yu M, Han J, et al. PEG-conjugated hemoglobin combination with cisplatin enforced the antiangiogenic effect in a cervical tumor xenograft model. *Artif Cells Blood Substit Immobil Biotechnol* 2008; 36: 487-97.
13. Efrati S, Berman S, Ben Aharon G, Siman-Tov Y, Averbukh Z, Weissgarten J. Application of normobaric hyperoxia therapy for amelioration of haemorrhagic shock-induced acute renal failure. *Nephrol Dial Transplant* 2008; 23: 2213-22.
14. Fan X, Krieg S, Kuo CJ, et al. VEGF blockade inhibits angiogenesis and reepithelialization of endometrium. *FASEB J* 2008.
15. Fujita N, Imai J, Suzuki T, et al. Vascular endothelial growth factor-A is a survival factor for nucleus pulposus cells in the intervertebral disc. *Biochem Biophys Res Commun* 2008; 372: 367-72.
16. Genin O, Hasdai A, Shinder D, Pines M. Hypoxia, hypoxia-inducible factor-1alpha (HIF-1alpha), and heat-shock proteins in tibial dyschondroplasia. *Poult Sci* 2008; 87: 1556-64.
17. Greijer AE, Delis-van Diemen PM, Fijneman RJ, et al. Presence of HIF-1 and related genes in normal mucosa, adenomas and carcinomas of the colorectum. *Virchows Arch* 2008; 452: 535-44.
18. Gulliksrud K, Vestvik IK, Galappathi K, Mathiesen B, Rofstad EK. Detection of different hypoxic cell subpopulations in human melanoma xenografts by pimonidazole immunohistochemistry. *Radiat Res* 2008; 170: 638-50.
19. Han J, Yu M, Dai M, et al. Effect of artificial oxygen carrier with chemotherapy on tumor hypoxia and neovascularization. *Artif Cells Blood Substit Immobil Biotechnol* 2008; 36: 431-8.
20. He F, Deng X, Wen B, et al. Noninvasive molecular imaging of hypoxia in human xenografts: comparing hypoxia-induced gene expression with endogenous and exogenous hypoxia markers. *Cancer Res* 2008; 68: 8597-606.

21. Heyman SN, Khamaisi M, Rosen S, Rosenberger C. Renal Parenchymal Hypoxia, Hypoxia Response and the Progression of Chronic Kidney Disease. *Am J Nephrol* 2008; 28: 998-1006.
22. Holcombe DJ, Lengefeld N, Gole GA, Barnett NL. The effects of acute intraocular pressure elevation on rat retinal glutamate transport. *Acta Ophthalmol* 2008; 86: 408-14.
23. Kalber TL, Waterton JC, Griffiths JR, Ryan AJ, Robinson SP. Longitudinal in vivo susceptibility contrast MRI measurements of LS174T colorectal liver metastasis in nude mice. *J Magn Reson Imaging* 2008; 28: 1451-8.
24. Klinkenberg LG, Sutherland LA, Bishai WR, Karakousis PC. Metronidazole Lacks Activity against Mycobacterium tuberculosis in an In Vivo Hypoxic Granuloma Model of Latency. *J Infect Dis* 2008; 198: 275-83.
25. Koch CJ. Importance of antibody concentration in the assessment of cellular hypoxia by flow cytometry: EF5 and pimonidazole. *Radiat Res* 2008; 169: 677-88
26. Kumar S, Brown SL, Kolozsvary A, Freytag SO, Kim JH. Efficacy of suicide gene therapy in hypoxic rat 9L glioma cells. *J Neurooncol* 2008.
27. Li X-F, JA OD. Hypoxia in microscopic tumors. *Cancer Letters* 2008; 264: 172-80.
28. Masamune A, Kikuta K, Watanabe T, Satoh K, Hirota M, Shimosegawa T. Hypoxia stimulates pancreatic stellate cells to induce fibrosis and angiogenesis in pancreatic cancer. *Am J Physiol Gastrointest Liver Physiol* 2008; 295: G709-17.
29. Mayr M, Sidibe A, Zampetaki A. The paradox of hypoxic and oxidative stress in atherosclerosis. *J Am Coll Cardiol* 2008; 51: 1266-7.
30. Miyazaki Y, Hara A, Kato K, et al. The effect of hypoxic microenvironment on matrix metalloproteinase expression in xenografts of human oral squamous cell carcinoma. *Int J Oncol* 2008; 32: 145-51.
31. Oh C, Dong Y, Harman C, Mighty HE, Kopelman J, Thompson LP. Chronic hypoxia differentially increases glutathione content and gamma-glutamyl cysteine synthetase expression in fetal guinea pig organs. *Early Hum Dev* 2008; 84: 121-7.
32. Peyssonnaud C, Boutin AT, Zinkernagel AS, Datta V, Nizet V, Johnson RS. Critical role of HIF-1alpha in keratinocyte defense against bacterial infection. *J Invest Dermatol* 2008; 128: 1964-8.
33. Rausch ME, Weisberg S, Vardhana P, Tortoriello DV. Obesity in C57BL/6J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration. *Int J Obes (Lond)* 2008; 32: 451-63.
34. Ream M, Ray AM, Chandra R, Chikaraishi DM. Early fetal hypoxia leads to growth restriction and myocardial thinning. *Am J Physiol Regul Integr Comp Physiol* 2008; 295: R583-95.
35. Ream MA, Ray AM, Chandra R, Chikaraishi DM. Early Fetal Hypoxia Leads to Growth Restriction and Myocardial Thinning. *Am J Physiol Regul Integr Comp Physiol* 2008.
36. Riedl CC, Brader P, Zanzonico PB, et al. Imaging hypoxia in orthotopic rat liver tumors with iodine 124-labeled iodoazomycin galactopyranoside PET. *Radiology* 2008; 248: 561-70.
37. Rosenberger C, Goldfarb M, Shina A, et al. Evidence for sustained renal hypoxia and transient hypoxia adaptation in experimental rhabdomyolysis-induced acute kidney injury. *Nephrol Dial Transplant* 2008; 23: 1135-43.
38. Rosenberger C, Khamaisi M, Abassi Z, et al. Adaptation to hypoxia in the diabetic rat kidney. *Kidney Int* 2008; 73: 34-42.

39. Rudolfsson SH, Bergh A. Testosterone-stimulated growth of the rat prostate may be driven by tissue hypoxia and hypoxia-inducible factor-1alpha. *J Endocrinol* 2008; 196: 11-9.
40. Saito Y, Uppal A, Byfield G, Budd S, Hartnett ME. Activated NAD(P)H Oxidase from Supplemental Oxygen Induces Neovascularization Independent of VEGF in Retinopathy of Prematurity Model. *Invest Ophthalmol Vis Sci* 2008; 49: 1591-8.
41. Scigliano S, Pinel S, Poussier S, et al. Measurement of hypoxia using invasive oxygen-sensitive electrode, pimonidazole binding and 18F-FDG uptake in anaemic or erythropoietin-treated mice bearing human glioma xenografts. *Int J Oncol* 2008; 32: 69-77.
42. Selvakumaran M, Yao KS, Feldman MD, O'Dwyer PJ. Antitumor effect of the angiogenesis inhibitor bevacizumab is dependent on susceptibility of tumors to hypoxia-induced apoptosis. *Biochem Pharmacol* 2008; 75: 627-38.
43. Sersa G, Jarm T, Kotnik T, et al. Vascular disrupting action of electroporation and electrochemotherapy with bleomycin in murine sarcoma. *Br J Cancer* 2008; 98: 388-98.
44. Sluimer JC, Gasc JM, van Wanroij JL, et al. Hypoxia, hypoxia-inducible transcription factor, and macrophages in human atherosclerotic plaques are correlated with intraplaque angiogenesis. *J Am Coll Cardiol* 2008; 51: 1258-65.
45. Sun X, Li XF, Russell J, et al. Changes in tumor hypoxia induced by mild temperature hyperthermia as assessed by dual-tracer immunohistochemistry. *Radiother Oncol* 2008.
46. Takahashi M, Yasui H, Ogura A, et al. X Irradiation Combined with TNF alpha-related Apoptosis-inducing Ligand (TRAIL) Reduces Hypoxic Regions of Human Gastric Adenocarcinoma Xenografts in SCID Mice. *J Radiat Res (Tokyo)* 2008; 49: 153-61.
47. Troost EG, Laverman P, Philippens ME, et al. Correlation of [(18)F]FMISO autoradiography and pimonidazole immunohistochemistry in human head and neck carcinoma xenografts. *Eur J Nucl Med Mol Imaging* 2008.
48. Troost EG, Laverman P, Philippens ME, et al. Correlation of [18F]FMISO autoradiography and pimonidazole [corrected] immunohistochemistry in human head and neck carcinoma xenografts. *Eur J Nucl Med Mol Imaging* 2008; 35: 1803-11.
49. Via LE, Lin PL, Ray SM, et al. Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and nonhuman primates. *Infect Immun* 2008; 76: 2333-40.
50. Vignozzi L, Morelli A, Filippi S, et al. Effect of sildenafil administration on penile hypoxia induced by cavernous neurotomy in the rat. *Int J Impot Res* 2008; 20: 60-7.
51. Wiggins KJ, Tiauw V, Zhang Y, Gilbert RE, Langham RG, Kelly DJ. Perindopril attenuates tubular hypoxia and inflammation in an experimental model of diabetic nephropathy in transgenic Ren-2 rats. *Nephrology (Carlton)* 2008; 13: 721-9.
52. Wijffels KI, Marres HA, Peters JP, Rijken PF, van der Kogel AJ, Kaanders JH. Tumour cell proliferation under hypoxic conditions in human head and neck squamous cell carcinomas. *Oral Oncol* 2008; 44: 335-44.
53. Yeom CJ, Chung JK, Kang JH, et al. Visualization of Hypoxia-Inducible Factor-1 Transcriptional Activation in C6 Glioma Using Luciferase and Sodium Iodide Symporter Genes. *J Nucl Med* 2008.
54. Yu M, Han J, Dai M, et al. Influence of PEG-conjugated hemoglobin on tumor oxygenation and response to chemotherapy. *Artif Cells Blood Substit Immobil Biotechnol* 2008; 36: 551-61.
55. Zhao D, Najbauer J, Garcia E, et al. Neural stem cell tropism to glioma: critical role of tumor hypoxia. *Mol Cancer Res* 2008; 6: 1819-29.

2007

1. Aly S, Laskay T, Mages J, Malzan A, Lang R, Ehlers S. Interferon-gamma-dependent mechanisms of mycobacteria-induced pulmonary immunopathology: the role of angiostasis and CXCR3-targeted chemokines for granuloma necrosis. *J Pathol* 2007; 212: 295-305.
2. Amarilio R, Viukov SV, Sharir A, Eshkar-Oren I, Johnson RS, Zelzer E. HIF1alpha regulation of Sox9 is necessary to maintain differentiation of hypoxic prechondrogenic cells during early skeletogenesis. *Development* 2007; 134: 3917-28.
3. Bernhardt WM, Wiesener MS, Weidemann A, et al. Involvement of hypoxia-inducible transcription factors in polycystic kidney disease. *Am J Pathol* 2007; 170: 830-42.
4. Brown DA, MacLellan WR, Laks H, Dunn JC, Wu BM, Beygui RE. Analysis of oxygen transport in a diffusion-limited model of engineered heart tissue. *Biotechnol Bioeng* 2007; 97: 962-75.
5. Burgu B, Medina Ortiz WE, Pitera JE, Woolf AS, Wilcox DT. Vascular endothelial growth factor mediates hypoxic stimulated embryonic bladder growth in organ culture. *J Urol* 2007; 177: 1552-7.
6. Bussink J, Kaanders JH, van der Kogel AJ. Microenvironmental transformations by VEGF- and EGF-receptor inhibition and potential implications for responsiveness to radiotherapy. *Radiother Oncol* 2007; 82: 10-7.
7. Chang YS, Adnane J, Trail PA, et al. Sorafenib (BAY 43-9006) inhibits tumor growth and vascularization and induces tumor apoptosis and hypoxia in RCC xenograft models. *Cancer Chemother Pharmacol* 2007; 59: 561-74.
8. Cheluvappa R, Hilmer SN, Kwun SY, Cogger VC, DG LEC. Effects of old age on hepatocyte oxygenation. *Ann N Y Acad Sci* 2007; 1114: 88-92.
9. Cheluvappa R, Hilmer SN, Kwun SY, et al. The effect of old age on liver oxygenation and the hepatic expression of VEGF and VEGFR2. *Exp Gerontol* 2007; 42: 1012-9.
10. Degrossoli A, Bosetto MC, Lima CB, Giorgio S. Expression of hypoxia-inducible factor 1alpha in mononuclear phagocytes infected with *Leishmania amazonensis*. *Immunol Lett* 2007; 114: 119-25.
11. Deng X, Luyendyk JP, Zou W, et al. Neutrophil interaction with the hemostatic system contributes to liver injury in rats cotreated with lipopolysaccharide and ranitidine. *J Pharmacol Exp Ther* 2007; 322: 852-61.
12. Donadieu E, Hamdi W, Deveze A, et al. Improved cryosections and specific immunohistochemical methods for detecting hypoxia in mouse and rat cochleae. *Acta Histochem* 2007; 109: 177-84.
13. Echevarria M, Munoz-Cabello AM, Sanchez-Silva R, Toledo-Aral JJ, Lopez-Barneo J. Development of cytosolic hypoxia and hypoxia-inducible factor stabilization are facilitated by aquaporin-1 expression. *J Biol Chem* 2007; 282: 30207-15.
14. Fang Y, Sullivan R, Graham CH. Confluence-dependent resistance to doxorubicin in human MDA-MB-231 breast carcinoma cells requires hypoxia-inducible factor-1 activity. *Exp Cell Res* 2007; 313: 867-77.
15. Frangoulis M, Georgiou P, Chrisostomidis C, et al. Rat epigastric flap survival and VEGF expression after local copper application. *Plast Reconstr Surg* 2007; 119: 837-43.

16. Gabellieri C, Eykyn TR, Leach MO. Conformational exchange in pimonidazole--a hypoxia marker. *Magn Reson Chem* 2007; 45: 621-3.
17. Gonzalez AM, Yazici I, Kusza K, Siemionow M. Effects of fresh versus banked blood transfusions on microcirculatory hemodynamics and tissue oxygenation in the rat cremaster model. *Surgery* 2007; 141: 630-9.
18. Higgins DF, Kimura K, Bernhardt WM, et al. Hypoxia promotes fibrogenesis in vivo via HIF-1 stimulation of epithelial-to-mesenchymal transition. *J Clin Invest* 2007; 117: 3810-20.
19. Hoogsteen IJ, Marres HA, Bussink J, van der Kogel AJ, Kaanders JH. Tumor microenvironment in head and neck squamous cell carcinomas: Predictive value and clinical relevance of hypoxic markers. A review. *Head Neck* 2007.
20. Hoskin PJ, Carnell DM, Taylor NJ, et al. Hypoxia in prostate cancer: correlation of BOLD-MRI with pimonidazole immunohistochemistry-initial observations. *Int J Radiat Oncol Biol Phys* 2007; 68: 1065-71.
21. Hosogai N, Fukuhara A, Oshima K, et al. Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes* 2007; 56: 901-11.
22. Iheagwara KN, Thom SR, Deutschman CS, Levy RJ. Myocardial cytochrome oxidase activity is decreased following carbon monoxide exposure. *Biochim Biophys Acta* 2007.
23. Imamura R, Moriyama T, Isaka Y, et al. Erythropoietin protects the kidneys against ischemia reperfusion injury by activating hypoxia inducible factor-1alpha. *Transplantation* 2007; 83: 1371-9.
24. Kaneko H, Yu D, Miura M. Overexpression of IGF-I receptor in HeLa cells enhances in vivo radioresponse. *Biochem Biophys Res Commun* 2007; 363: 937-41.
25. Karantza-Wadsworth V, Patel S, Kravchuk O, et al. Autophagy mitigates metabolic stress and genome damage in mammary tumorigenesis. *Genes Dev* 2007; 21: 1621-35.
26. Kim YJ, Kang HH, Ahn JH, Chung JW. Hypoxic changes in the central nervous system of noise-exposed mice. *Acta Otolaryngol Suppl* 2007: 73-7.
27. Krause M, Prager J, Zhou X, et al. EGFR-TK inhibition before radiotherapy reduces tumour volume but does not improve local control: differential response of cancer stem cells and nontumorigenic cells? *Radiother Oncol* 2007; 83: 316-25.
28. Le QT. Identifying and targeting hypoxia in head and neck cancer: a brief overview of current approaches. *Int J Radiat Oncol Biol Phys* 2007; 69: S56-8.
29. Lenaerts AJ, Hoff D, Aly S, et al. Location of persisting mycobacteria in a Guinea pig model of tuberculosis revealed by r207910. *Antimicrob Agents Chemother* 2007; 51: 3338-45.
30. Levesque JP, Winkler IG, Hendy J, et al. Hematopoietic Progenitor Cell Mobilization Results in Hypoxia with Increased Hypoxia-Inducible Transcription Factor-1[alpha] and Vascular Endothelial Growth Factor A in Bone Marrow. *Stem Cells* 2007; 25: 1954-65.
31. Li XF, Carlin S, Urano M, Russell J, Ling CC, O'Donoghue JA. Visualization of hypoxia in microscopic tumors by immunofluorescent microscopy. *Cancer Res* 2007; 67: 7646-53.
32. Ljungkvist AS, Bussink J, Kaanders JH, van der Kogel AJ. Dynamics of tumor hypoxia measured with bioreductive hypoxic cell markers. *Radiat Res* 2007; 167: 127-45.
33. Matsumoto K, Szajek L, Krishna MC, et al. The influence of tumor oxygenation on hypoxia imaging in murine squamous cell carcinoma using [64Cu]Cu-ATSM or [18F]Fluoromisonidazole positron emission tomography. *Int J Oncol* 2007; 30: 873-81.

34. McClelland MR, Carskadon SL, Zhao L, et al. Diversity of the angiogenic phenotype in non-small cell lung cancer. *Am J Respir Cell Mol Biol* 2007; 36: 343-50.
35. Noto T, Furuichi Y, Ishiye M, et al. Tacrolimus (FK506) limits accumulation of granulocytes and platelets and protects against brain damage after transient focal cerebral ischemia in rat. *Biol Pharm Bull* 2007; 30: 313-7.
36. Olmsted-Davis E, Gannon FH, Ozen M, et al. Hypoxic adipocytes pattern early heterotopic bone formation. *Am J Pathol* 2007; 170: 620-32.
37. Papadopoulou MV, Bloomer WD, Taylor AP, Hernandez M, Blumenthal RD, Hollingshead MG. Advantage of combining NLCQ-1 (NSC 709257) with radiation in treatment of human head and neck xenografts. *Radiat Res* 2007; 168: 65-71.
38. Parmar K, Mauch P, Vergilio JA, Sackstein R, Down JD. Distribution of hematopoietic stem cells in the bone marrow according to regional hypoxia. *Proc Natl Acad Sci U S A* 2007; 104: 5431-6.
39. Plock J, Frese S, Keogh A, et al. Activation of non-ischemic, hypoxia-inducible signalling pathways up-regulate cytoprotective genes in the murine liver. *J Hepatol* 2007; 47: 538-45.
40. Pringle KG, Kind KL, Thompson JG, Roberts CT. Complex interactions between hypoxia inducible factors, insulin-like growth factor-II and oxygen in early murine trophoblasts. *Placenta* 2007; 28: 1147-57.
41. Riva C, Donadieu E, Magnan J, Lavieille JP. Age-related hearing loss in CD/1 mice is associated to ROS formation and HIF target proteins up-regulation in the cochlea. *Exp Gerontol* 2007; 42: 327-36.
42. Rofstad EK, Galappathi K, Mathiesen B, Ruud EB. Fluctuating and diffusion-limited hypoxia in hypoxia-induced metastasis. *Clin Cancer Res* 2007; 13: 1971-8.
43. Shafer J, Davis AR, Gannon FH, et al. Oxygen tension directs chondrogenic differentiation of myelo-monocytic progenitors during endochondral bone formation. *Tissue Eng* 2007; 13: 2011-9.
44. Shahrzad S, Bertrand K, Minhas K, Coomber BL. Induction of DNA hypomethylation by tumor hypoxia. *Epigenetics* 2007; 2: 119-25.
45. Shin KH, Diaz-Gonzalez JA, Russell J, et al. Detecting changes in tumor hypoxia with carbonic anhydrase IX and pimonidazole. *Cancer Biol Ther* 2007; 6: 70-5.
46. Silva P, Homer JJ, Slevin NJ, et al. Clinical and biological factors affecting response to radiotherapy in patients with head and neck cancer: a review. *Clin Otolaryngol* 2007; 32: 337-45.
47. Terada N, Ohno N, Saitoh S, Ohno S. Immunohistochemical detection of hypoxia in mouse liver tissues treated with pimonidazole using "in vivo cryotechnique". *Histochem Cell Biol* 2007; 128: 253-61.
48. Thored P, Wood J, Arvidsson A, Cammenga J, Kokaia Z, Lindvall O. Long-Term Neuroblast Migration Along Blood Vessels in an Area With Transient Angiogenesis and Increased Vascularization After Stroke. *Stroke* 2007; 38: 3032-9.
49. Tugues S, Fernandez-Varo G, Munoz-Luque J, et al. Antiangiogenic treatment with sunitinib ameliorates inflammatory infiltrate, fibrosis, and portal pressure in cirrhotic rats. *Hepatology* 2007; 46: 1919-26.
50. van der Bilt JD, Soeters ME, Duyverman AM, et al. Perinecrotic hypoxia contributes to ischemia/reperfusion-accelerated outgrowth of colorectal micrometastases. *Am J Pathol* 2007; 170: 1379-88.

51. Westbury CB, Pearson A, Nerurkar A, et al. Hypoxia can be detected in irradiated normal human tissue: a study using the hypoxic marker pimonidazole hydrochloride. *Br J Radiol* 2007; 80: 934-8.
52. Wijffels KI, Marres HA, Peters JP, Rijken PF, van der Kogel AJ, Kaanders JH. Tumour cell proliferation under hypoxic conditions in human head and neck squamous cell carcinomas. *Oral Oncol* 2007.
53. Ye J, Gao Z, Yin J, He Q. Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice. *Am J Physiol Endocrinol Metab* 2007; 293: E1118-28.
54. Zips D, Yaromina A, Schutze C, et al. Experimental evaluation of functional imaging for radiotherapy. *Strahlenther Onkol* 2007; 183 Spec No 2: 41-2.

2006

1. Aly S, Wagner K, Keller C, et al. Oxygen status of lung granulomas in *Mycobacterium tuberculosis*-infected mice. *J Pathol* 2006; 210: 298-305.
2. Badger WJ, Whitbeck C, Kogan B, Chichester P, Levin RM. The immediate effect of castration on female rabbit bladder blood flow and tissue oxygenation. *Urol Int* 2006; 76: 264-8.
3. Carnell DM, Smith RE, Daley FM, Saunders MI, Bentzen SM, Hoskin PJ. An immunohistochemical assessment of hypoxia in prostate carcinoma using pimonidazole: Implications for radioresistance. *Int J Radiat Oncol Biol Phys* 2006; 65: 91-9.
4. Ceelen W, Smeets P, Backes W, et al. Noninvasive monitoring of radiotherapy-induced microvascular changes using dynamic contrast enhanced magnetic resonance imaging (DCE-MRI) in a colorectal tumor model. *Int J Radiat Oncol Biol Phys* 2006; 64: 1188-96.
5. Cheema AN, Hong T, Nili N, et al. Adventitial microvessel formation after coronary stenting and the effects of SU11218, a tyrosine kinase inhibitor. *J Am Coll Cardiol* 2006; 47: 1067-75.
6. Copple BL, Roth RA, Ganey PE. Anticoagulation and inhibition of nitric oxide synthase influence hepatic hypoxia after monocrotaline exposure. *Toxicology* 2006; 225: 128-37.
7. Danylkova NO, Pomeranz HD, Alcalá SR, McLoon LK. Histological and morphometric evaluation of transient retinal and optic nerve ischemia in rat. *Brain Res* 2006; 1096: 20-9.
8. de Gooyer TE, Stevenson KA, Humphries P, et al. Rod photoreceptor loss in Rho-/- mice reduces retinal hypoxia and hypoxia-regulated gene expression. *Invest Ophthalmol Vis Sci* 2006; 47: 5553-60.
9. de Gooyer TE, Stevenson KA, Humphries P, Simpson DA, Gardiner TA, Stitt AW. Retinopathy is reduced during experimental diabetes in a mouse model of outer retinal degeneration. *Invest Ophthalmol Vis Sci* 2006; 47: 5561-8.
10. De Schutter H, Barbe B, Spaepen M, et al. Microsatellite alterations in head and neck squamous cell carcinoma and relation to expression of pimonidazole, CA IX and GLUT-1. *Radiother Oncol* 2006; 80: 143-50.
11. De Schutter HD, Barbe B, Spaepen M, et al. Microsatellite alterations in head and neck squamous cell carcinoma and relation to expression of pimonidazole, CA IX and GLUT-1. *Radiother Oncol* 2006; 80: 143-50.

12. Dorow DS, Cullinane C, Conus N, et al. Multi-tracer small animal PET imaging of the tumour response to the novel pan-Erb-B inhibitor CI-1033. *Eur J Nucl Med Mol Imaging* 2006; 1-12.
13. Durand RE, Aquino-Parsons C. The fate of hypoxic (pimonidazole-labelled) cells in human cervix tumours undergoing chemo-radiotherapy. *Radiother Oncol* 2006; 80: 138-42.
14. Franco M, Man S, Chen L, et al. Targeted anti-vascular endothelial growth factor receptor-2 therapy leads to short-term and long-term impairment of vascular function and increase in tumor hypoxia. *Cancer Res* 2006; 66: 3639-48.
15. Goethals L, Debuquoy A, Perneel C, et al. Hypoxia in human colorectal adenocarcinoma: Comparison between extrinsic and potential intrinsic hypoxia markers. *Int J Radiat Oncol Biol Phys* 2006; 65: 246-54.
16. Goldfarb M, Rosenberger C, Abassi Z, et al. Acute-on-chronic renal failure in the rat: functional compensation and hypoxia tolerance. *Am J Nephrol* 2006; 26: 22-33.
17. Gookin JL, Copple CN, Papich MG, et al. Efficacy of ronidazole for treatment of feline *Tritrichomonas foetus* infection. *J Vet Intern Med* 2006; 20: 536-43.
18. Greco O, Joiner MC, Doleh A, Powell AD, Hillman GG, Scott SD. Hypoxia- and radiation-activated Cre/loxP 'molecular switch' vectors for gene therapy of cancer. *Gene Ther* 2006; 13: 206-15.
19. Hammarsten P, Halin S, Wikstom P, Henriksson R, Rudolfsson SH, Bergh A. Inhibitory effects of castration in an orthotopic model of androgen-independent prostate cancer can be mimicked and enhanced by angiogenesis inhibition. *Clin Cancer Res* 2006; 12: 7431-6.
20. Huxham LA, Kyle AH, Baker JH, McNicol KL, Minchinton AI. Tirapazamine causes vascular dysfunction in HCT-116 tumour xenografts. *Radiother Oncol* 2006; 78: 138-45.
21. Jankovic B, Aquino-Parsons C, Raleigh JA, et al. Comparison between pimonidazole binding, oxygen electrode measurements, and expression of endogenous hypoxia markers in cancer of the uterine cervix. *Cytometry B Clin Cytom* 2006; 70: 45-55.
22. Jensen RL. Hypoxia in the tumorigenesis of gliomas and as a potential target for therapeutic measures. *Neurosurg Focus* 2006; 20: E24.
23. Jonathan RA, Wijffels KI, Peeters W, et al. The prognostic value of endogenous hypoxia-related markers for head and neck squamous cell carcinomas treated with ARCON. *Radiother Oncol* 2006; 79: 288-97.
24. Kabuubi P, Loncaster JA, Davidson SE, et al. No relationship between thymidine phosphorylase (TP, PD-ECGF) expression and hypoxia in carcinoma of the cervix. *Br J Cancer* 2006; 94: 115-20.
25. Kang SH, Cho HT, Devi S, et al. Antitumor effect of 2-methoxyestradiol in a rat orthotopic brain tumor model. *Cancer Res* 2006; 66: 11991-7.
26. Khan Z, Michalopoulos GK, Stolz DB. Peroxisomal localization of hypoxia-inducible factors and hypoxia-inducible factor regulatory hydroxylases in primary rat hepatocytes exposed to hypoxia-reoxygenation. *Am J Pathol* 2006; 169: 1251-69.
27. Kirkpatrick JP, Hardee ME, Snyder SA, et al. The effect of darbepoetin alfa on growth, oxygenation and radioresponsiveness of a breast adenocarcinoma. *Radiat Res* 2006; 165: 192-201.

28. Kleiter MM, Thrall DE, Malarkey DE, et al. A comparison of oral and intravenous pimonidazole in canine tumors using intravenous CCI-103F as a control hypoxia marker. *Int J Radiat Oncol Biol Phys* 2006; 64: 592-602.
29. Laderoute KR, Amin K, Calaoagan JM, et al. 5'-AMP-activated protein kinase (AMPK) is induced by low-oxygen and glucose deprivation conditions found in solid-tumor microenvironments. *Mol Cell Biol* 2006; 26: 5336-47.
30. Lipnik K, Greco O, Scott S, et al. Hypoxia- and radiation-inducible, breast cell-specific targeting of retroviral vectors. *Virology* 2006; 349: 121-33.
31. Ljungkvist AS, Bussink J, Kaanders JH, Wiedenmann NE, Vlasman R, van der Kogel AJ. Dynamics of hypoxia, proliferation and apoptosis after irradiation in a murine tumor model. *Radiat Res* 2006; 165: 326-36.
32. Lokmic Z, Darby IA, Thompson EW, Mitchell GM. Time course analysis of hypoxia, granulation tissue and blood vessel growth, and remodeling in healing rat cutaneous incisional primary intention wounds. *Wound Repair Regen* 2006; 14: 277-88.
33. Minchinton AI, Tannock IF. Drug penetration in solid tumours. *Nat Rev Cancer* 2006; 6: 583-92.
34. Morani A, Barros RP, Imamov O, et al. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci U S A* 2006; 103: 7165-9.
35. Nanka O, Valasek P, Dvorakova M, Grim M. Experimental hypoxia and embryonic angiogenesis. *Dev Dyn* 2006; 235: 723-33.
36. Nordmark M, Loncaster J, Aquino-Parsons C, et al. The prognostic value of pimonidazole and tumour pO₂ in human cervix carcinomas after radiation therapy: A prospective international multi-center study. *Radiother Oncol* 2006; 80: 123-31.
37. Noto T, Furuichi Y, Ishiye M, et al. Temporal and topographic profiles of tissue hypoxia following transient focal cerebral ischemia in rats. *J Vet Med Sci* 2006; 68: 803-7.
38. Papandreou I, Cairns RA, Fontana L, Lim AL, Denko NC. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab* 2006; 3: 187-97.
39. Pugachev A, Carlin S, Ling C, Humm J. SU-FF-J-89: Intratumoral Pattern of FDG Uptake in Human Xenograft Models. *Medical Physics* 2006; 33: 2040-.
40. Raleigh JA, Arcasoy MO, Amikn K, Lininger RA, Varia MA. Comparison between exogenous and endogenous hypoxia markers in human cancer: pimonidazole and erythropoietin. . American Association for Cancer Research 97th Annual Meeting; 2006; Washington, DC; 2006. p. Abstract #3631.
41. Raleigh JA, Lee DY-W, Ji X-S, inventors; Natural Pharmacia International, Incorporated, assignee. Fluorinated 2-nitroimidazoles for the non-invasive detection of tissue hypoxia. USA. 2006.
42. Rosenberger C, Rosen S, Shina A, et al. Hypoxia-inducible factors and tubular cell survival in isolated perfused kidneys. *Kidney Int* 2006; 70: 60-70.
43. Safran M, Kim WY, O'Connell F, et al. Mouse model for noninvasive imaging of HIF prolyl hydroxylase activity: assessment of an oral agent that stimulates erythropoietin production. *Proc Natl Acad Sci U S A* 2006; 103: 105-10.
44. Schaffer CB, Friedman B, Nishimura N, et al. Two-photon imaging of cortical surface microvessels reveals a robust redistribution in blood flow after vascular occlusion. *PLoS Biol* 2006; 4: e22.

45. Stockwin LH, Blonder J, Bumke MA, et al. Proteomic analysis of plasma membrane from hypoxia-adapted malignant melanoma. *J Proteome Res* 2006; 5: 2996-3007.
46. Tanaka H, Yamamoto M, Hashimoto N, et al. Hypoxia-independent overexpression of hypoxia-inducible factor 1alpha as an early change in mouse hepatocarcinogenesis. *Cancer Res* 2006; 66: 11263-70.
47. Tanaka T, Kato H, Kojima I, et al. Hypoxia and expression of hypoxia-inducible factor in the aging kidney. *J Gerontol A Biol Sci Med Sci* 2006; 61: 795-805.
48. Thorwarth D, Eschmann SM, Holzner F, Paulsen F, Alber M. Combined uptake of [18F]FDG and [18F]FMISO correlates with radiation therapy outcome in head-and-neck cancer patients. *Radiother Oncol* 2006; 80: 151-6.
49. Troost EG, Laverman P, Kaanders JH, et al. Imaging hypoxia after oxygenation-modification: Comparing [(18)F]FMISO autoradiography with pimonidazole immunohistochemistry in human xenograft tumors. *Radiother Oncol* 2006.
50. van Laarhoven HW, Kaanders JH, Lok J, et al. Hypoxia in relation to vasculature and proliferation in liver metastases in patients with colorectal cancer. *Int J Radiat Oncol Biol Phys* 2006; 64: 473-82.
51. Vignozzi L, Filippi S, Morelli A, et al. Effect of chronic tadalafil administration on penile hypoxia induced by cavernous neurotomy in the rat. *J Sex Med* 2006; 3: 419-31.
52. Villanueva S, Cespedes C, Vio CP. Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. *Am J Physiol Regul Integr Comp Physiol* 2006; 290: R861-70.
53. Webster WS, Howe AM, Abela D, Oakes DJ. The relationship between cleft lip, maxillary hypoplasia, hypoxia and phenytoin. *Curr Pharm Des* 2006; 12: 1431-48.
54. Yaromina A, Zips D, Thames HD, et al. Pimonidazole labelling and response to fractionated irradiation of five human squamous cell carcinoma (hSCC) lines in nude mice: The need for a multivariate approach in biomarker studies. *Radiother Oncol* 2006; 81: 122-9.
55. Yuan H, Schroeder T, Bowsher JE, Hedlund LW, Wong T, Dewhirst MW. Intertumoral differences in hypoxia selectivity of the PET imaging agent ⁶⁴Cu(II)-diacetyl-bis(N4-methylthiosemicarbazone). *J Nucl Med* 2006; 47: 989-98.
56. Zampino M, Yuzhakova M, Hansen J, et al. Sex-related dimorphic response of HIF-1 alpha expression in myocardial ischemia. *Am J Physiol Heart Circ Physiol* 2006; 291: H957-64.

2005

1. Airley RE, Phillips RM, Evans AE, et al. Hypoxia-regulated glucose transporter Glut-1 may influence chemosensitivity to some alkylating agents: Results of EORTC (First Translational Award) study of the relevance of tumour hypoxia to the outcome of chemotherapy in human tumour-derived xenografts. *Int J Oncol* 2005; 26: 1477-84.
2. Arcasoy MO, Amin K, Chou SC, Haroon ZA, Varia M, Raleigh JA. Erythropoietin and erythropoietin receptor expression in head and neck cancer: relationship to tumor hypoxia. *Clin Cancer Res* 2005; 11: 20-7.
3. Asosingh K, De Raeve H, de Ridder M, et al. Role of the hypoxic bone marrow microenvironment in 5T2MM murine myeloma tumor progression. *Haematologica* 2005; 90: 810-7.

4. Cao Y, Li CY, Moeller BJ, et al. Observation of incipient tumor angiogenesis that is independent of hypoxia and hypoxia inducible factor-1 activation. *Cancer Res* 2005; 65: 5498-505.
5. Casanovas O, Hicklin DJ, Bergers G, Hanahan D. Drug resistance by evasion of antiangiogenic targeting of VEGF signaling in late-stage pancreatic islet tumors. *Cancer Cell* 2005; 8: 299-309.
6. Covington MD, Bayless KJ, Burghardt RC, Davis GE, Parrish AR. Ischemia-induced cleavage of cadherins in NRK cells: evidence for a role of metalloproteinases. *Am J Physiol Renal Physiol* 2005; 289: F280-8.
7. Damaser MS, Whitbeck C, Chichester P, Levin RM. Effect of vaginal distension on blood flow and hypoxia of urogenital organs of the female rat. *J Appl Physiol* 2005; 98: 1884-90.
8. Danielsson BR, Johansson A, Danielsson C, Azarbayjani F, Blomgren B, Skold AC. Phenytoin teratogenicity: Hypoxia marker and effects on embryonic heart rhythm suggest an hERG-related mechanism. *Birth Defects Res A Clin Mol Teratol* 2005; 73: 146-53.
9. Doege K, Heine S, Jensen I, Jelkmann W, Metzner E. Inhibition of mitochondrial respiration elevates oxygen concentration but leaves regulation of hypoxia-inducible factor (HIF) intact. *Blood* 2005; 106: 2311-7.
10. Forder JP, Munzenmaier DH, Greene AS. Angiogenic protection from focal ischemia with angiotensin II type 1 receptor blockade in the rat. *Am J Physiol Heart Circ Physiol* 2005; 288: H1989-96.
11. Gaber T, Dziurla R, Tripmacher R, Burmester GR, Buttgereit F. Hypoxia inducible factor (HIF) in rheumatology: low O₂! See what HIF can do! *Ann Rheum Dis* 2005; 64: 971-80.
12. Gardiner TA, Gibson DS, de Gooyer TE, de la Cruz VF, McDonald DM, Stitt AW. Inhibition of Tumor Necrosis Factor-[alpha] Improves Physiological Angiogenesis and Reduces Pathological Neovascularization in Ischemic Retinopathy. *Am J Pathol* 2005; 166: 637-44.
13. Harada H, Kizaka-Kondoh S, Hiraoka M. Optical imaging of tumor hypoxia and evaluation of efficacy of a hypoxia-targeting drug in living animals. *Mol Imaging* 2005; 4: 182-93.
14. Hofer SO, Mitchell GM, Penington AJ, et al. The use of pimonidazole to characterise hypoxia in the internal environment of an in vivo tissue engineering chamber. *Br J Plast Surg* 2005; 58: 1104-14.
15. Hoogsteen IJ, Peeters WJ, Marres HA, et al. Erythropoietin receptor is not a surrogate marker for tumor hypoxia and does not correlate with survival in head and neck squamous cell carcinomas. *Radiother Oncol* 2005; 76: 213-8.
16. Janssen HL, Haustermans KM, Balm AJ, Begg AC. Hypoxia in head and neck cancer: how much, how important? *Head Neck* 2005; 27: 622-38.
17. Kempf VA, Lebedziejewski M, Alitalo K, et al. Activation of hypoxia-inducible factor-1 in bacillary angiomatosis: evidence for a role of hypoxia-inducible factor-1 in bacterial infections. *Circulation* 2005; 111: 1054-62.
18. Lewis C, Murdoch C. Macrophage responses to hypoxia: implications for tumor progression and anti-cancer therapies. *Am J Pathol* 2005; 167: 627-35.
19. Lin AD, Mannikarottu A, Chaudhry A, et al. Protective effects of grape suspension on in vivo ischaemia/reperfusion of the rabbit bladder. *BJU Int* 2005; 96: 1397-402.

20. Liu J, Qu R, Ogura M, Shibata T, Harada H, Hiraoka M. Real-time Imaging of Hypoxia-inducible Factor-1 Activity in Tumor Xenografts. *J Radiat Res (Tokyo)* 2005; 46: 93-102.
21. Ljungkvist AS, Bussink J, Kaanders JH, et al. Hypoxic cell turnover in different solid tumor lines. *Int J Radiat Oncol Biol Phys* 2005; 62: 1157-68.
22. Lunt SJ, Telfer BA, Fitzmaurice RJ, Stratford IJ, Williams KJ. Tirapazamine administered as a neoadjuvant to radiotherapy reduces metastatic dissemination. *Clin Cancer Res* 2005; 11: 4212-6.
23. Luyendyk JP, Shaw PJ, Green CD, Maddox JF, Ganey PE, Roth RA. Coagulation-mediated hypoxia and neutrophil-dependent hepatic injury in rats given lipopolysaccharide and ranitidine. *J Pharmacol Exp Ther* 2005; 314: 1023-31.
24. Mayer A, Hockel M, Vaupel P. Carbonic anhydrase IX expression and tumor oxygenation status do not correlate at the microregional level in locally advanced cancers of the uterine cervix. *Clin Cancer Res* 2005; 11: 7220-5.
25. Mizukami Y, Jo WS, Duerr EM, et al. Induction of interleukin-8 preserves the angiogenic response in HIF-1alpha-deficient colon cancer cells. *Nat Med* 2005; 11: 992-7.
26. Murdoch C, Lewis CE. Macrophage migration and gene expression in response to tumor hypoxia. *Int J Cancer* 2005; 117: 701-8.
27. Nelson DW, Cao H, Zhu Y, et al. A noninvasive approach for assessing tumor hypoxia in xenografts: developing a urinary marker for hypoxia. *Cancer Res* 2005; 65: 6151-8.
28. O'Donoghue J A, Zanzonico P, Pugachev A, et al. Assessment of regional tumor hypoxia using (18)F-fluoromisonidazole and (64)Cu(II)-diacetyl-bis(N4-methylthiosemicarbazone) positron emission tomography: Comparative study featuring microPET imaging, Po(2) probe measurement, autoradiography, and fluorescent microscopy in the R3327-AT and FaDu rat tumor models. *Int J Radiat Oncol Biol Phys* 2005; 61: 1493-502.
29. Paizis G, Tikellis C, Cooper ME, et al. Chronic liver injury in rats and humans upregulates the novel enzyme angiotensin converting enzyme 2. *Gut* 2005; 54: 1790-6.
30. Pollard M, Suckow MA. Hormone-refractory prostate cancer in the Lobund-Wistar rat. *Exp Biol Med (Maywood)* 2005; 230: 520-6.
31. Pugachev A, Ruan S, Carlin S, et al. Dependence of FDG uptake on tumor microenvironment. *Int J Radiat Oncol Biol Phys* 2005; 62: 545-53.
32. Rofstad EK, Mathiesen B, Henriksen K, Kindem K, Galappathi K. The Tumor Bed Effect: increased metastatic dissemination from hypoxia-Induced up-regulation of metastasis-promoting gene products. *Cancer Res* 2005; 65: 2387-96.
33. Rosenberger C, Heyman SN, Rosen S, et al. Up-regulation of HIF in experimental acute renal failure: Evidence for a protective transcriptional response to hypoxia. *Kidney Int* 2005; 67: 531-42.
34. Schuurin J, Bussink J, Bernsen HJ, Peeters W, van Der Kogel AJ. Irradiation combined with SU5416: microvascular changes and growth delay in a human xenograft glioblastoma tumor line. *Int J Radiat Oncol Biol Phys* 2005; 61: 529-34.
35. Sobhanifar S, Aquino-Parsons C, Stanbridge EJ, Olive P. Reduced Expression of Hypoxia-Inducible Factor-1[alpha] in Perinecrotic Regions of Solid Tumors. *Cancer Res* 2005; 65: 7259-66.
36. Solomon B, Binns D, Roselt P, et al. Modulation of intratumoral hypoxia by the epidermal growth factor receptor inhibitor gefitinib detected using small animal PET imaging. *Mol Cancer Ther* 2005; 4: 1417-22.

37. Thorwarth D, Eschmann SM, Paulsen F, Alber M. A kinetic model for dynamic [(18)F]-Fmiso PET data to analyse tumour hypoxia. *Phys Med Biol* 2005; 50: 2209-24.
38. Troost EG, Bussink J, Kaanders JH, et al. Comparison of different methods of CAIX quantification in relation to hypoxia in three human head and neck tumor lines. *Radiother Oncol* 2005; 76: 194-9.
39. van der Schaft DWJ, Hillen F, Pauwels P, et al. Tumor Cell Plasticity in Ewing Sarcoma, an Alternative Circulatory System Stimulated by Hypoxia. *Cancer Res* 2005; 65: 11520-8.
40. van Herpen CM, Bussink J, van der Kogel AJ, et al. Interleukin-12 has no effect on vascular density, perfusion, hypoxia and proliferation of an implanted human squamous cell carcinoma xenograft tumour despite up-regulation of ICAM-1. *Anticancer Res* 2005; 25: 1015-10121.
41. van Laarhoven HW, de Geus-Oei LF, Wiering B, et al. Gadopentetate dimeglumine and FDG uptake in liver metastases of colorectal carcinoma as determined with MR imaging and PET. *Radiology* 2005; 237: 181-8.
42. Villanueva S, Cespedes C, Vio CP. Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. *Am J Physiol Regul Integr Comp Physiol* 2005.
43. Vordermark D, Kraft P, Katzer A, Bolling T, Willner J, Flentje M. Glucose requirement for hypoxic accumulation of hypoxia-inducible factor-1alpha (HIF-1alpha). *Cancer Lett* 2005; 230: 122-33.
44. Wang PX, Sanders PW. Mechanism of hypertensive nephropathy in the Dahl/Rapp rat: a primary disorder of vascular smooth muscle. *Am J Physiol Renal Physiol* 2005; 288: F236-42.
45. Williams KJ, Parker CA, Stratford IJ. Exogenous and endogenous markers of tumour oxygenation status: definitive markers of tumour hypoxia? *Adv Exp Med Biol* 2005; 566: 285-94.
46. Yahiro T, Masui S, Kubota N, Yamada K, Kobayashi A, Kishii K. Effects of hypoxic cell radiosensitizer doranidazole (PR-350) on the radioresponse of murine and human tumor cells in vitro and in vivo. *J Radiat Res (Tokyo)* 2005; 46: 363-72.
46. Yaromina A, Holscher T, Eicheler W, et al. Does heterogeneity of pimonidazole labelling correspond to the heterogeneity of radiation-response of FaDu human squamous cell carcinoma? *Radiother Oncol* 2005; 76: 206-12.

2004

1. Bhattacharya A, Toth K, Mazurchuk R, et al. Lack of microvessels in well-differentiated regions of human head and neck squamous cell carcinoma A253 associated with functional magnetic resonance imaging detectable hypoxia, limited drug delivery, and resistance to irinotecan therapy. *Clin Cancer Res* 2004; 10: 8005-17.
2. Buchler P, Reber HA, Lavey RS, et al. Tumor hypoxia correlates with metastatic tumor growth of pancreatic cancer in an orthotopic murine model. *J Surg Res* 2004; 120: 295-303.
3. Cardenas-Navia LI, Yu D, Braun RD, Brizel DM, Secomb TW, Dewhirst MW. Tumor-dependent kinetics of partial pressure of oxygen fluctuations during air and oxygen breathing. *Cancer Res* 2004; 64: 6010-7.

4. Chou SC, Azuma Y, Varia MA, Raleigh JA. Evidence that involucrin, a marker for differentiation, is oxygen regulated in human squamous cell carcinomas. *Br J Cancer* 2004; 90: 728-35.
5. Copple BL, Rondelli CM, Maddox JF, Hoglen NC, Ganey PE, Roth RA. Modes of cell death in rat liver after monocrotaline exposure. *Toxicol Sci* 2004; 77: 172-82.
6. Davies Cd, Lundstrom LM, Frengen J, et al. Radiation improves the distribution and uptake of liposomal doxorubicin (caelyx) in human osteosarcoma xenografts. *Cancer Res* 2004; 64: 547-53.
7. Dearling JL, Flynn AA, Sutcliffe-Goulden J, et al. Analysis of the regional uptake of radiolabeled deoxyglucose analogs in human tumor xenografts. *J Nucl Med* 2004; 45: 101-7.
8. Dubois L, Landuyt W, Haustermans K, et al. Evaluation of hypoxia in an experimental rat tumour model by [(18)F]fluoromisonidazole PET and immunohistochemistry. *Br J Cancer* 2004; 91: 1947-54.
9. Erler JT, Cawthorne CJ, Williams KJ, et al. Hypoxia-mediated down-regulation of Bid and Bax in tumors occurs via hypoxia-inducible factor 1-dependent and -independent mechanisms and contributes to drug resistance. *Mol Cell Biol* 2004; 24: 2875-89.
10. Fei P, Wang W, Kim SH, et al. Bnip3L is induced by p53 under hypoxia, and its knockdown promotes tumor growth. *Cancer Cell* 2004; 6: 597-609.
11. Freeburg PB, Abrahamson DR. Divergent expression patterns for hypoxia-inducible factor-1beta and aryl hydrocarbon receptor nuclear transporter-2 in developing kidney. *J Am Soc Nephrol* 2004; 15: 2569-78.
12. Hoskin PJ, Sibtain A, Daley FM, Saunders MI, Wilson GD. The immunohistochemical assessment of hypoxia, vascularity and proliferation in bladder carcinoma. *Radiother Oncol* 2004; 72: 159-68.
13. Huang J, Frischer JS, New T, et al. TNP-470 promotes initial vascular sprouting in xenograft tumors. *Mol Cancer Ther* 2004; 3: 335-43.
14. Hutchison GJ, Valentine HR, Loncaster JA, et al. Hypoxia-inducible factor 1alpha expression as an intrinsic marker of hypoxia: correlation with tumor oxygen, pimonidazole measurements, and outcome in locally advanced carcinoma of the cervix. *Clin Cancer Res* 2004; 10: 8405-12.
15. Huxham LA, Kyle AH, Baker JH, Nykilchuk LK, Minchinton AI. Microregional effects of gemcitabine in HCT-116 xenografts. *Cancer Res* 2004; 64: 6537-41.
16. Janssen HL, Hoebbers FJ, Sprong D, et al. Differentiation-associated staining with anti-pimonidazole antibodies in head and neck tumors. *Radiother Oncol* 2004; 70: 91-7.
17. Jeong WI, Do SH, Yun HS, et al. Hypoxia potentiates transforming growth factor-beta expression of hepatocyte during the cirrhotic condition in rat liver. *Liver Int* 2004; 24: 658-68.
18. Kostourou V, Troy H, Murray JF, et al. Overexpression of dimethylarginine dimethylaminohydrolase enhances tumor hypoxia: an insight into the relationship of hypoxia and angiogenesis in vivo. *Neoplasia* 2004; 6: 401-11.
19. Levin R, Chichester P, Levin S, Buttyan R. Role of angiogenesis in bladder response to partial outlet obstruction. *Scand J Urol Nephrol Suppl* 2004: 37-47.
20. Manotham K, Tanaka T, Matsumoto M, et al. Evidence of tubular hypoxia in the early phase in the remnant kidney model. *J Am Soc Nephrol* 2004; 15: 1277-88.

21. Matsumoto M, Tanaka T, Yamamoto T, et al. Hypoperfusion of Peritubular Capillaries Induces Chronic Hypoxia before Progression of Tubulointerstitial Injury in a Progressive Model of Rat Glomerulonephritis. *J Am Soc Nephrol* 2004; 15: 1574-81.
22. Murdoch C, Giannoudis A, Lewis CE. Mechanisms regulating the recruitment of macrophages into hypoxic areas of tumors and other ischemic tissues. *Blood* 2004; 104: 2224-34.
23. Peters CL, Morris CJ, Mapp PI, Blake DR, Lewis CE, Winrow VR. The transcription factors hypoxia-inducible factor 1alpha and Ets-1 colocalize in the hypoxic synovium of inflamed joints in adjuvant-induced arthritis. *Arthritis Rheum* 2004; 50: 291-6.
24. Post DE, Devi NS, Li Z, et al. Cancer therapy with a replicating oncolytic adenovirus targeting the hypoxic microenvironment of tumors. *Clin Cancer Res* 2004; 10: 8603-12.
25. Prabhakaran K, Sampson DA, Hoehner JC. Neuroblastoma survival and death: an in vitro model of hypoxia and metabolic stress. *J Surg Res* 2004; 116: 288-96.
26. Rofstad EK, Mathiesen B, Galappathi K. Increased metastatic dissemination in human melanoma xenografts after subcurative radiation treatment: radiation-induced increase in fraction of hypoxic cells and hypoxia-induced up-regulation of urokinase-type plasminogen activator receptor. *Cancer Res* 2004; 64: 13-8.
27. Samoszuk MK, Walter J, Mechetner E. Improved immunohistochemical method for detecting hypoxia gradients in mouse tissues and tumors. *J Histochem Cytochem* 2004; 52: 837-9.
28. Tang N, Wang L, Esko J, et al. Loss of HIF-1alpha in endothelial cells disrupts a hypoxia-driven VEGF autocrine loop necessary for tumorigenesis. *Cancer Cell* 2004; 6: 485-95.
29. van Laarhoven H, Lok J, Rijpkema M, et al. Relation between dynamic gadolinium uptake rate, tumor vasculature and tumor hypoxia in human colorectal liver metastases. *Proc Intl Soc Mag Reson Med*; 2004; 2004. p. 148.
30. van Laarhoven HW, Bussink J, Lok J, Punt CJ, Heerschap A, van Der Kogel AJ. Effects of nicotinamide and carbogen in different murine colon carcinomas: immunohistochemical analysis of vascular architecture and microenvironmental parameters. *Int J Radiat Oncol Biol Phys* 2004; 60: 310-21.
31. van Wijngaarden J, de Rooij K, van Beek E, et al. Identification of differentially expressed genes in a renal cell carcinoma tumor model after endostatin-treatment. *Lab Invest* 2004; 84: 1472-83.
32. Yang L, DeBusk LM, Fukuda K, et al. Expansion of myeloid immune suppressor Gr⁺CD11b⁺ cells in tumor-bearing host directly promotes tumor angiogenesis. *Cancer Cell* 2004; 6: 409-21.
33. Zips D, Adam M, Flentje M, et al. Impact of Hypoxia and the Metabolic Microenvironment on Radiotherapy of Solid Tumors Introduction of a Multiinstitutional Research Project. *Strahlenther Onkol* 2004; 180: 609-15.

2003

1. Airley RE, Loncaster J, Raleigh JA, et al. GLUT-1 and CAIX as intrinsic markers of hypoxia in carcinoma of the cervix: Relationship to pimonidazole binding. *Int J Cancer* 2003; 104: 85-91.

2. Azuma Y, Chou SC, Lininger RA, Murphy BJ, Varia MA, Raleigh JA. Hypoxia and differentiation in squamous cell carcinomas of the uterine cervix: pimonidazole and involucrin. *Clin Cancer Res* 2003; 9: 4944-452.
3. Basile DP, Donohoe DL, Roethe K, Mattson DL. Chronic renal hypoxia after acute ischemic injury: effects of L-arginine on hypoxia and secondary damage. *Am J Physiol Renal Physiol* 2003; 284: F338-48.
4. Bassnett S, McNulty R. The effect of elevated intraocular oxygen on organelle degradation in the embryonic chicken lens. *J Exp Biol* 2003; 206: 4353-61.
5. Begg AC. Is HIF-1alpha a good marker for tumor hypoxia? *Int J Radiat Oncol Biol Phys* 2003; 56: 917-9.
6. Blouw B, Song H, Tihan T, et al. The hypoxic response of tumors is dependent on their microenvironment. *Cancer Cell* 2003; 4: 133-46.
7. Bussink J, Kaanders JH, van der Kogel AJ. Tumor hypoxia at the micro-regional level: clinical relevance and predictive value of exogenous and endogenous hypoxic cell markers. *Radiother Oncol* 2003; 67: 3-15.
8. Cachat F, Lange-Sperandio B, Chang AY, et al. Ureteral obstruction in neonatal mice elicits segment-specific tubular cell responses leading to nephron loss. *Kidney Int* 2003; 63: 564-75.
9. Choy MY, Leung TN, Leung PS, Lau TK. First Trimester Human Trophoblast Production Of Placental Corticotrophin-Releasing Hormone (CRH) Is Unresponsive To Hypoxia In-Vitro. *Early Pregnancy: Biology and Medicine* 2003; VI: 235-47.
10. Danielsson BR, Skold AC, Johansson A, Dillner B, Blomgren B. Teratogenicity by the hERG potassium channel blocking drug almokalant: use of hypoxia marker gives evidence for a hypoxia-related mechanism mediated via embryonic arrhythmia. *Toxicol Appl Pharmacol* 2003; 193: 168-76.
11. Freeburg PB, Robert B, St John PL, Abrahamson DR. Podocyte expression of hypoxia-inducible factor (HIF)-1 and HIF-2 during glomerular development. *J Am Soc Nephrol* 2003; 14: 927-38.
12. Hoskin PJ, Sibtain A, Daley FM, Wilson GD. GLUT1 and CAIX as intrinsic markers of hypoxia in bladder cancer: relationship with vascularity and proliferation as predictors of outcome of ARCON. *Br J Cancer* 2003; 89: 1290-7.
13. Levin RM, O'Connor LJ, Leggett RE, Whitbeck C, Chichester P. Focal hypoxia of the obstructed rabbit bladder wall correlates with intermediate decompensation. *Neurourol Urodyn* 2003; 22: 156-63.
14. McKim SE, Uesugi T, Raleigh JA, McClain CJ, Arteel GE. Chronic intragastric alcohol exposure causes hypoxia and oxidative stress in the rat pancreas. *Arch Biochem Biophys* 2003; 417: 34-43.
15. Nordmark M, Loncaster J, Aquino-Parsons C, et al. Measurements of hypoxia using pimonidazole and polarographic oxygen-sensitive electrodes in human cervix carcinomas. *Radiother Oncol* 2003; 67: 35-44.
16. Petersen C, Eicheler W, Frommel A, et al. Proliferation and micromilieu during fractionated irradiation of human FaDu squamous cell carcinoma in nude mice. *Int J Radiat Biol* 2003; 79: 469-77.
17. Samoszuk M, Corwin MA. Mast cell inhibitor cromolyn increases blood clotting and hypoxia in murine breast cancer. *Int J Cancer* 2003; 107: 159-63.

18. Sminia P, Acker H, Eikesdal HP, et al. Oxygenation and response to irradiation of organotypic multicellular spheroids of human glioma. *Anticancer Res* 2003; 23: 1461-6.
19. Vordermark D, Brown JM. Endogenous markers of tumor hypoxia predictors of clinical radiation resistance? *Strahlenther Onkol* 2003; 179: 801-11.
20. Vordermark D, Brown JM. Evaluation of hypoxia-inducible factor-1alpha (HIF-1alpha) as an intrinsic marker of tumor hypoxia in U87 MG human glioblastoma: in vitro and xenograft studies. *Int J Radiat Oncol Biol Phys* 2003; 56: 1184-93.
21. Zhao D, Ran S, Constantinescu A, Hahn EW, Mason RP. Tumor oxygen dynamics: correlation of in vivo MRI with histological findings. *Neoplasia* 2003; 5: 308-18.
22. Zoula S, Rijken PF, Peters JP, et al. Pimonidazole binding in C6 rat brain glioma: relation with lipid droplet detection. *Br J Cancer* 2003; 88: 1439-44.

2002

1. Arcasoy MO, Amin K, Karayal AF, et al. Functional significance of erythropoietin receptor expression in breast cancer. *Lab Invest* 2002; 82: 911-8.
2. Bardag-Gorce F, French BA, Li J, et al. The importance of cycling of blood alcohol levels in the pathogenesis of experimental alcoholic liver disease in rats. *Gastroenterology* 2002; 123: 325-35.
3. Bennewith KL, Raleigh JA, Durand RE. Orally administered pimonidazole to label hypoxic tumor cells. *Cancer Res* 2002; 62: 6827-30.
4. Corpechot C, Barbu V, Wendum D, et al. Hepatocyte growth factor and c-Met inhibition by hepatic cell hypoxia: a potential mechanism for liver regeneration failure in experimental cirrhosis. *Am J Pathol* 2002; 160: 613-20.
5. Ghafar MA, Anastasiadis AG, Olsson LE, et al. Hypoxia and an angiogenic response in the partially obstructed rat bladder. *Lab Invest* 2002; 82: 903-9.
6. Ghafar MA, Shabsigh A, Chichester P, et al. Effects of chronic partial outlet obstruction on blood flow and oxygenation of the rat bladder. *J Urol* 2002; 167: 1508-12.
7. Hale LP, Braun RD, Gwinn WM, Greer PK, Dewhirst MW. Hypoxia in the thymus: role of oxygen tension in thymocyte survival. *Am J Physiol Heart Circ Physiol* 2002; 282: H1467-77.
8. Harada H, Hiraoka M, Kizaka-Kondoh S. Antitumor effect of TAT-oxygen-dependent degradation-caspase-3 fusion protein specifically stabilized and activated in hypoxic tumor cells. *Cancer Res* 2002; 62: 2013-8.
9. Hoebbers FJ, Janssen HL, Olmos AV, et al. Phase 1 study to identify tumour hypoxia in patients with head and neck cancer using technetium-99m BRU 59-21. Vascular architecture, hypoxia, and proliferation in first-generation xenografts of human head-and-neck squamous cell carcinomas. Why do hypoxic cells behave badly? *Eur J Nucl Med Mol Imaging* 2002; 29: 1206-11.
10. Janssen HL, Haustermans KM, Sprong D, et al. HIF-1alpha, pimonidazole, and iododeoxyuridine to estimate hypoxia and perfusion in human head-and-neck tumors. *Int J Radiat Oncol Biol Phys* 2002; 54: 1537-49.
11. Kaanders JH, Wijffels KI, Marres HA, et al. Pimonidazole binding and tumor vascularity predict for treatment outcome in head and neck cancer. *Cancer Res* 2002; 62: 7066-74.

12. Olive PL, Banath JP, Durand RE. The range of oxygenation in SiHa tumor xenografts. *Radiat Res* 2002; 158: 159-66.
13. Olive PL, Luo CM, Banath JP. Local hypoxia is produced at sites of intratumour injection. *Br J Cancer* 2002; 86: 429-35.
14. Pedley RB, El-Emir Eb, Flynn AA, et al. Synergy between vascular targeting agents and antibody-directed therapy. *Int J Radiat Oncol Biol Phys* 2002; 54: 1524-31.
15. Rijken PF, Peters JP, Van der Kogel AJ. Quantitative analysis of varying profiles of hypoxia in relation to functional vessels in different human glioma xenograft lines. *Radiat Res* 2002; 157: 626-32.
16. Rofstad EK, Halsor EF. Hypoxia-associated spontaneous pulmonary metastasis in human melanoma xenografts: involvement of microvascular hot spots induced in hypoxic foci by interleukin 8. *Br J Cancer* 2002; 86: 301-8.
17. Rofstad EK, Rasmussen H, Galappathi K, Mathiesen B, Nilsen K, Graff BA. Hypoxia promotes lymph node metastasis in human melanoma xenografts by up-regulating the urokinase-type plasminogen activator receptor. *Cancer Res* 2002; 62: 1847-53.
18. Rofstad EK, Tunheim SH, Mathiesen B, et al. Pulmonary and lymph node metastasis is associated with primary tumor interstitial fluid pressure in human melanoma xenografts. *Cancer Res* 2002; 62: 661-4.
19. Yin M, Zhong Z, Connor HD, et al. Protective effect of glycine on renal injury induced by ischemia- reperfusion in vivo. *Am J Physiol Renal Physiol* 2002; 282: F417-23.

2000

1. Bernsen HJ, Rijken PF, Peters H, et al. Hypoxia in a human intracerebral glioma model. *J Neurosurg* 2000; 93: 449-54.
2. Bonn D. Why do hypoxic cells behave badly? *Lancet Oncol* 2000; 1: 202.
3. Bussink J, Kaanders JH, Rijken PF, Raleigh JA, Van der Kogel AJ. Changes in blood perfusion and hypoxia after irradiation of a human squamous cell carcinoma xenograft tumor line. *Radiat Res* 2000; 153: 398-404.
4. Bussink J, Kaanders JH, Strik AM, van der Kogel AJ. Effects of nicotinamide and carbogen on oxygenation in human tumor xenografts measured with luminescence based fiber-optic probes. *Radiother Oncol* 2000; 57: 21-30.
5. Bussink J, Kaanders JH, Strik AM, Vojnovic B, van Der Kogel AJ. Optical sensor-based oxygen tension measurements correspond with hypoxia marker binding in three human tumor xenograft lines. *Radiat Res* 2000; 154: 547-55.
6. Danielsen T, Rofstad EK. The constitutive level of vascular endothelial growth factor (VEGF) is more important than hypoxia-induced VEGF up-regulation in the angiogenesis of human melanoma xenografts. *Br J Cancer* 2000; 82: 1528-34.
7. Haroon ZA, Raleigh JA, Greenberg CS, Dewhirst MW. Early wound healing exhibits cytokine surge without evidence of hypoxia. *Ann Surg* 2000; 231: 137-47.
8. Haustermans K, Hofland I, Van de Pavert L, et al. Diffusion limited hypoxia estimated by vascular image analysis: comparison with pimonidazole staining in human tumors. *Radiother Oncol* 2000; 55: 325-33.
9. Lehnert S. Prediction of tumor response to therapy: molecular markers and the microenvironment. Apoptosis and chips: an overview of the proceedings. *Radiat Res* 2000; 154: 121-4.

10. Ljungkvist AS, Bussink J, Rijken PF, Raleigh JA, Denekamp J, Van Der Kogel AJ. Changes in tumor hypoxia measured with a double hypoxic marker technique. *Int J Radiat Oncol Biol Phys* 2000; 48: 1529-38.
11. Olive PL, Durand RE, Raleigh JA, Luo C, Aquino-Parsons C. Comparison between the comet assay and pimonidazole binding for measuring tumour hypoxia. *Br J Cancer* 2000; 83: 1525-31.
12. Raleigh JA, Chou SC, Calkins-Adams DP, Ballenger CA, Novotny DB, Varia MA. A clinical study of hypoxia and metallothionein protein expression in squamous cell carcinomas. *Clin Cancer Res* 2000; 6: 855-62.
13. Rijken PF, Bernsen HJ, Peters JP, Hodgkiss RJ, Raleigh JA, van der Kogel AJ. Spatial relationship between hypoxia and the (perfused) vascular network in a human glioma xenograft: a quantitative multi-parameter analysis. *Int J Radiat Oncol Biol Phys* 2000; 48: 571-82.
14. Thurman RG. Sex-related liver injury due to alcohol involves activation of Kupffer cells by endotoxin. *Can J Gastroenterol* 2000; 14 Suppl D: 129D-35D.
15. Wijffels KI, Kaanders JH, Rijken PF, et al. Vascular architecture and hypoxic profiles in human head and neck squamous cell carcinomas. *Br J Cancer* 2000; 83: 674-83.
16. Wykoff CC, Beasley NJ, Watson PH, et al. Hypoxia-inducible expression of tumor-associated carbonic anhydrases. *Cancer Res* 2000; 60: 7075-83.

1980-1990s

1. Laurent F, Benard P, Canal P, Soula G. Autoradiographic distribution of [¹⁴C]-labelled pimonidazole in rhabdomyosarcoma-bearing rats and pigmented mice. *Cancer Chemother Pharmacol* 1988; 22: 308-15.
2. Cobb LM, Nolan J, Butler SA. Distribution of pimonidazole and RSU 1069 in tumour and normal tissues. *Br J Cancer* 1990; 62: 915-8.
3. Arteel GE, Thurman RG, Yates JM, Raleigh JA. Evidence that hypoxia markers detect oxygen gradients in liver: pimonidazole and retrograde perfusion of rat liver. *Br J Cancer* 1995; 72: 889-95.
4. Arteel GE, Raleigh JA, Bradford BU, Thurman RG. Acute alcohol produces hypoxia directly in rat liver tissue in vivo: role of Kupffer cells. *Am J Physiol* 1996; 271: G494-500.
5. Chou SC, Flood PM, Raleigh JA. Marking hypoxic cells for complement and cytotoxic T lymphocyte-mediated lysis: using pimonidazole. *Br J Cancer Suppl* 1996; 27: S213-6.
6. Cowan DS, Hicks KO, Wilson WR. Multicellular membranes as an in vitro model for extravascular diffusion in tumours. *Br J Cancer Suppl* 1996; 27: S28-31.
7. Raleigh JA, Dewhirst MW, Thrall DE. Measuring tumor hypoxia. *Semin Radiat Oncol* 1996; 6: 37-45.
8. Arteel GE, Iimuro Y, Yin M, Raleigh JA, Thurman RG. Chronic enteral ethanol treatment causes hypoxia in rat liver tissue in vivo. *Hepatology* 1997; 25: 920-6.
9. Azuma C, Raleigh JA, Thrall DE. Longevity of pimonidazole adducts in spontaneous canine tumors as an estimate of hypoxic cell lifetime. *Radiat Res* 1997; 148: 35-42.
10. Kennedy AS, Raleigh JA, Perez GM, et al. Proliferation and hypoxia in human squamous cell carcinoma of the cervix: first report of combined immunohistochemical assays. *Int J Radiat Oncol Biol Phys* 1997; 37: 897-905.

11. Lyng H, Sundfor K, Rofstad EK. Oxygen tension in human tumours measured with polarographic needle electrodes and its relationship to vascular density, necrosis and hypoxia. *Radiother Oncol* 1997; 44: 163-9.
12. Raleigh JA, Lee DY, inventors; Raleigh, J.A. Lee, D.Y., assignee. Derivatives of 2-nitroimidazoles as hypoxic cell markers. USA patent 5.674,693. 1997 October 7, 1997.
13. Stachlewitz RF, Arteel GE, Raleigh JA, Connor HD, Mason RP, Thurman RG. Development and characterization of a new model of tacrine-induced hepatotoxicity: role of the sympathetic nervous system and hypoxia- reoxygenation. *J Pharmacol Exp Ther* 1997; 282: 1591-9.
14. Tamulevicius P, Streffer C. Bioluminescence imaging of metabolites in a human tumour xenograft after treatment with hyperthermia and/or the radiosensitizer pimonidazole. *Int J Hyperthermia* 1997; 13: 235-45.
15. Arteel GE, Thurman RG, Raleigh JA. Reductive metabolism of the hypoxia marker pimonidazole is regulated by oxygen tension independent of the pyridine nucleotide redox state. *Eur J Biochem* 1998; 253: 743-50.
16. Durand RE, Raleigh JA. Identification of nonproliferating but viable hypoxic tumor cells in vivo. *Cancer Res* 1998; 58: 3547-50.
17. Raleigh JA, Calkins-Adams DP, Rinker LH, et al. Hypoxia and vascular endothelial growth factor expression in human squamous cell carcinomas using pimonidazole as a hypoxia marker. *Cancer Res* 1998; 58: 3765-8.
18. Raleigh JA, Chou SC, Tables L, Suchindran S, Varia MA, Horsman MR. Relationship of hypoxia to metallothionein expression in murine tumors. *Int J Radiat Oncol Biol Phys* 1998; 42: 727-30.
19. Thurman RG. II. Alcoholic liver injury involves activation of Kupffer cells by endotoxin. *Am J Physiol* 1998; 275: G605-11.
20. Thurman RG, Bradford BU, Iimuro Y, et al. The role of gut-derived bacterial toxins and free radicals in alcohol- induced liver injury. *J Gastroenterol Hepatol* 1998; 13 Suppl: S39-50.
21. Varia MA, Calkins-Adams DP, Rinker LH, et al. Pimonidazole: a novel hypoxia marker for complementary study of tumor hypoxia and cell proliferation in cervical carcinoma. *Gynecol Oncol* 1998; 71: 270-7.
22. Zhong Z, Arteel GE, Connor HD, et al. Cyclosporin A increases hypoxia and free radical production in rat kidneys: prevention by dietary glycine. *Am J Physiol* 1998; 275: F595-604.
23. Bernsen HJ, Rijken PF, Peters JP, et al. Suramin treatment of human glioma xenografts; effects on tumor vasculature and oxygenation status. *J Neurooncol* 1999; 44: 129-36.
24. Dodd JS, Raleigh JA, Gross TS. Osteocyte hypoxia: a novel mechanotransduction pathway. *Am J Physiol Renal Physiol* 1999; 277: C598-602.
25. Raleigh JA, Chou SC, Arteel GE, Horsman MR. Comparisons among pimonidazole binding, oxygen electrode measurements, and radiation response in C3H mouse tumors. *Radiat Res* 1999; 151: 580-9.
26. Rofstad EK, Maseide K. Radiobiological and immunohistochemical assessment of hypoxia in human melanoma xenografts: acute and chronic hypoxia in individual tumours. *Int J Radiat Biol* 1999; 75: 1377-93.

27. Rosmorduc O, Wendum D, Corpechot C, et al. Hepatocellular hypoxia-induced vascular endothelial growth factor expression and angiogenesis in experimental biliary cirrhosis. *Am J Pathol* 1999; 155: 1065-73.
28. Schemmer P, Bradford BU, Rose ML, et al. Intravenous glycine improves survival in rat liver transplantation. *Am J Physiol* 1999; 276: G924-32.
29. Schemmer P, Bunzendahl H, Raleigh JA, Thurman RG. Graft survival is improved by hepatic denervation before organ harvesting. *Transplantation* 1999; 67: 1301-7.
30. Schemmer P, Connor HD, Arteel GE, et al. Reperfusion injury in livers due to gentle in situ organ manipulation during harvest involves hypoxia and free radicals. *J Pharmacol Exp Ther* 1999; 290: 235-40.
31. Sinusas AJ. The potential for myocardial imaging with hypoxia markers. *Semin Nucl Med* 1999; 29: 330-8.
32. Zhong Z, Arteel GE, Connor HD, et al. Binge drinking disturbs hepatic microcirculation after transplantation: prevention with free radical scavengers. *J Pharmacol Exp Ther* 1999; 290: 611-20.