

## All References to Hypoxyprobe-1 Use in Chronological Order.

1. Arteel, G. E., Thurman, R. G., Yates, J. M., and Raleigh, J. A. Evidence that hypoxia markers detect oxygen gradients in liver: pimonidazole and retrograde perfusion of rat liver. *Br J Cancer*, 72: 889-895, 1995.
2. Arteel, G. E., Raleigh, J. A., Bradford, B. U., and Thurman, R. G. Acute alcohol produces hypoxia directly in rat liver tissue in vivo: role of Kupffer cells. *Am J Physiol*, 271: G494-500, 1996.
3. Chou, S. C., Flood, P. M., and Raleigh, J. A. Marking hypoxic cells for complement and cytotoxic T lymphocyte-mediated lysis: using pimonidazole. *Br J Cancer Suppl*, 27: S213-216, 1996.
4. Cowan, D. S., Hicks, K. O., and Wilson, W. R. Multicellular membranes as an in vitro model for extravascular diffusion in tumours. *Br J Cancer Suppl*, 27: S28-31, 1996.
5. Arteel, G. E., Iimuro, Y., Yin, M., Raleigh, J. A., and Thurman, R. G. Chronic enteral ethanol treatment causes hypoxia in rat liver tissue in vivo. *Hepatology*, 25: 920-926, 1997.
6. Azuma, C., Raleigh, J. A., and Thrall, D. E. Longevity of pimonidazole adducts in spontaneous canine tumors as an estimate of hypoxic cell lifetime. *Radiat Res*, 148: 35-42, 1997.
7. Kennedy, A. S., Raleigh, J. A., Perez, G. M., Calkins, D. P., Thrall, D. E., Novotny, D. B., and Varia, M. A. Proliferation and hypoxia in human squamous cell carcinoma of the cervix: first report of combined immunohistochemical assays. *Int J Radiat Oncol Biol Phys*, 37: 897-905, 1997.
8. Lyng, H., Sundfor, K., and Rofstad, E. K. Oxygen tension in human tumours measured with polarographic needle electrodes and its relationship to vascular density, necrosis and hypoxia. *Radiother Oncol*, 44: 163-169, 1997.
9. Stachlewitz, R. F., Arteel, G. E., Raleigh, J. A., Connor, H. D., Mason, R. P., and Thurman, R. G. Development and characterization of a new model of tacrine-induced hepatotoxicity: role of the sympathetic nervous system and hypoxia-reoxygenation. *J Pharmacol Exp Ther*, 282: 1591-1599, 1997.
10. Tamulevicius, P. and Streffer, C. Bioluminescence imaging of metabolites in a human tumour xenograft after treatment with hyperthermia and/or the radiosensitizer pimonidazole. *Int J Hyperthermia*, 13: 235-245, 1997.
11. Arteel, G. E., Thurman, R. G., and Raleigh, J. A. Reductive metabolism of the hypoxia marker pimonidazole is regulated by oxygen tension independent of the pyridine nucleotide redox state. *Eur J Biochem*, 253: 743-750, 1998.
12. Durand, R. E. and Raleigh, J. A. Identification of nonproliferating but viable hypoxic tumor cells in vivo. *Cancer Res*, 58: 3547-3550, 1998.
13. Raleigh, J. A., Calkins-Adams, D. P., Rinker, L. H., Ballenger, C. A., Weissler, M. C., Fowler, W. C., Jr., Novotny, D. B., and Varia, M. A. Hypoxia and vascular endothelial growth factor expression in human squamous cell carcinomas using pimonidazole as a hypoxia marker. *Cancer Res*, 58: 3765-3768, 1998.

14. Raleigh, J. A., Chou, S. C., Tables, L., Suchindran, S., Varia, M. A., and Horsman, M. R. Relationship of hypoxia to metallothionein expression in murine tumors. *Int J Radiat Oncol Biol Phys*, *42*: 727-730, 1998.
15. Thurman, R. G. II. Alcoholic liver injury involves activation of Kupffer cells by endotoxin. *Am J Physiol*, *275*: G605-611, 1998.
16. Thurman, R. G., Bradford, B. U., Iimuro, Y., Knecht, K. T., Arteel, G. E., Yin, M., Connor, H. D., Wall, C., Raleigh, J. A., Frankenberg, M. V., Adachi, Y., Forman, D. T., Brenner, D., Kadiiska, M., and Mason, R. P. The role of gut-derived bacterial toxins and free radicals in alcohol-induced liver injury. *J Gastroenterol Hepatol*, *13 Suppl*: S39-50, 1998.
17. Varia, M. A., Calkins-Adams, D. P., Rinker, L. H., Kennedy, A. S., Novotny, D. B., Fowler, W. C., Jr., and Raleigh, J. A. Pimonidazole: a novel hypoxia marker for complementary study of tumor hypoxia and cell proliferation in cervical carcinoma. *Gynecol Oncol*, *71*: 270-277, 1998.
18. Zhong, Z., Arteel, G. E., Connor, H. D., Yin, M., Frankenberg, M. V., Stachlewitz, R. F., Raleigh, J. A., Mason, R. P., and Thurman, R. G. Cyclosporin A increases hypoxia and free radical production in rat kidneys: prevention by dietary glycine. *Am J Physiol*, *275*: F595-604, 1998.
19. Bernsen, H. J., Rijken, P. F., Peters, J. P., Bakker, J. H., Boerman, R. H., Wesseling, P., and van der Kogel, A. J. Suramin treatment of human glioma xenografts; effects on tumor vasculature and oxygenation status. *J Neurooncol*, *44*: 129-136, 1999.
20. Dodd, J. S., Raleigh, J. A., and Gross, T. S. Osteocyte hypoxia: a novel mechanotransduction pathway. *Am. J. Physiol. Renal. Physiol.*, *277*: C598-602, 1999.
21. Raleigh, J. A., Chou, S. C., Arteel, G. E., and Horsman, M. R. Comparisons among pimonidazole binding, oxygen electrode measurements, and radiation response in C3H mouse tumors. *Radiat Res*, *151*: 580-589, 1999.
22. Rofstad, E. K. and Maseide, K. Radiobiological and immunohistochemical assessment of hypoxia in human melanoma xenografts: acute and chronic hypoxia in individual tumours. *Int J Radiat Biol*, *75*: 1377-1393, 1999.
23. Rosmorduc, O., Wendum, D., Corpechot, C., Galy, B., Sebbagh, N., Raleigh, J., Housset, C., and Poupon, R. Hepatocellular hypoxia-induced vascular endothelial growth factor expression and angiogenesis in experimental biliary cirrhosis. *Am J Pathol*, *155*: 1065-1073, 1999.
24. Schemmer, P., Bradford, B. U., Rose, M. L., Bunzendahl, H., Raleigh, J. A., Lemasters, J. J., and Thurman, R. G. Intravenous glycine improves survival in rat liver transplantation. *Am J Physiol*, *276*: G924-932, 1999.
25. Schemmer, P., Bunzendahl, H., Raleigh, J. A., and Thurman, R. G. Graft survival is improved by hepatic denervation before organ harvesting. *Transplantation*, *67*: 1301-1307, 1999.
26. Schemmer, P., Connor, H. D., Arteel, G. E., Raleigh, J. A., Bunzendahl, H., Mason, R. P., and Thurman, R. G. Reperfusion injury in livers due to gentle in situ organ manipulation during harvest involves hypoxia and free radicals. *J Pharmacol Exp Ther*, *290*: 235-240, 1999.

27. Sinusas, A. J. The potential for myocardial imaging with hypoxia markers. *Semin Nucl Med*, 29: 330-338, 1999.
28. Zhong, Z., Arteel, G. E., Connor, H. D., Schemmer, P., Chou, S. C., Raleigh, J. A., Mason, R. P., Lemasters, J. J., and Thurman, R. G. Binge drinking disturbs hepatic microcirculation after transplantation: prevention with free radical scavengers. *J Pharmacol Exp Ther*, 290: 611-620, 1999.
29. Bernsen, H. J., Rijken, P. F., Peters, H., Raleigh, J. A., Jeuken, J. W., Wesseling, P., and van der Kogel, A. J. Hypoxia in a human intracerebral glioma model. *J Neurosurg*, 93: 449-454, 2000.
30. Bonn, D. Why do hypoxic cells behave badly? *Lancet Oncol*, 1: 202, 2000.
31. Bussink, J., Kaanders, J. H., Rijken, P. F., Raleigh, J. A., and Van der Kogel, A. J. Changes in blood perfusion and hypoxia after irradiation of a human squamous cell carcinoma xenograft tumor line. *Radiat Res*, 153: 398-404, 2000.
32. Bussink, J., Kaanders, J. H., Strik, A. M., and van der Kogel, A. J. Effects of nicotinamide and carbogen on oxygenation in human tumor xenografts measured with luminescence based fiber-optic probes. *Radiother Oncol*, 57: 21-30, 2000.
33. Bussink, J., Kaanders, J. H., Strik, A. M., Vojnovic, B., and van Der Kogel, A. J. Optical sensor-based oxygen tension measurements correspond with hypoxia marker binding in three human tumor xenograft lines. *Radiat Res*, 154: 547-555, 2000.
34. Danielsen, T. and Rofstad, E. K. The constitutive level of vascular endothelial growth factor (VEGF) is more important than hypoxia-induced VEGF up-regulation in the angiogenesis of human melanoma xenografts. *Br J Cancer*, 82: 1528-1534, 2000.
35. Haroon, Z. A., Raleigh, J. A., Greenberg, C. S., and Dewhirst, M. W. Early wound healing exhibits cytokine surge without evidence of hypoxia. *Ann Surg*, 231: 137-147, 2000.
36. Haustermans, K., Hofland, I., Van de Pavert, L., Geboes, K., Varia, M., Raleigh, J., and Begg, A. C. Diffusion limited hypoxia estimated by vascular image analysis: comparison with pimonidazole staining in human tumors. *Radiother Oncol*, 55: 325-333, 2000.
37. Lehnert, S. Prediction of tumor response to therapy: molecular markers and the microenvironment. Apoptosis and chips: an overview of the proceedings. *Radiat Res*, 154: 121-124, 2000.
38. Ljungkvist, A. S., Bussink, J., Rijken, P. F., Raleigh, J. A., Denekamp, J., and Van Der Kogel, A. J. Changes in tumor hypoxia measured with a double hypoxic marker technique. *Int J Radiat Oncol Biol Phys*, 48: 1529-1538, 2000.
39. Olive, P. L., Durand, R. E., Raleigh, J. A., Luo, C., and Aquino-Parsons, C. Comparison between the comet assay and pimonidazole binding for measuring tumour hypoxia. *Br J Cancer*, 83: 1525-1531, 2000.
40. Raleigh, J. A., Chou, S. C., Calkins-Adams, D. P., Ballenger, C. A., Novotny, D. B., and Varia, M. A. A clinical study of hypoxia and metallothionein protein expression in squamous cell carcinomas. *Clin Cancer Res*, 6: 855-862, 2000.
41. Rijken, P. F., Bernsen, H. J., Peters, J. P., Hodgkiss, R. J., Raleigh, J. A., and van der Kogel, A. J. Spatial relationship between hypoxia and the (perfused) vascular

- network in a human glioma xenograft: a quantitative multi-parameter analysis. *Int J Radiat Oncol Biol Phys*, 48: 571-582, 2000.
42. Thurman, R. G. Sex-related liver injury due to alcohol involves activation of Kupffer cells by endotoxin. *Can J Gastroenterol*, 14 Suppl D: 129D-135D, 2000.
  43. Wijffels, K. I., Kaanders, J. H., Rijken, P. F., Bussink, J., van den Hoogen, F. J., Marres, H. A., de Wilde, P. C., Raleigh, J. A., and van der Kogel, A. J. Vascular architecture and hypoxic profiles in human head and neck squamous cell carcinomas. *Br J Cancer*, 83: 674-683, 2000.
  44. Wykoff, C. C., Beasley, N. J., Watson, P. H., Turner, K. J., Pastorek, J., Sibtain, A., Wilson, G. D., Turley, H., Talks, K. L., Maxwell, P. H., Pugh, C. W., Ratcliffe, P. J., and Harris, A. L. Hypoxia-inducible expression of tumor-associated carbonic anhydrases. *Cancer Res*, 60: 7075-7083, 2000.
  45. Albina, J. E., Mastrofrancesco, B., Vessella, J. A., Louis, C. A., Henry, W. L., Jr., and Reichner, J. S. HIF-1 expression in healing wounds: HIF-1alpha induction in primary inflammatory cells by TNF-alpha. *Am J Physiol Cell Physiol*, 281: C1971-1977, 2001.
  46. Begg, A. C., Janssen, H., Sprong, D., Hofland, I., Blommestijn, G., Raleigh, J. A., Varia, M., Balm, A., Van Velthuyzen, L., Delaere, P., Sciot, R., and Haustermans, K. M. G. Hypoxia and perfusion measurements in human tumors--initial experience with pimonidazole and IUdR. *Acta Oncol*, 40: 924-928, 2001.
  47. hbEikesdal, H. P., Bjerkvig, R., Raleigh, J. A., Mella, O., and Dahl, O. Tumor vasculature is targeted by the combination of combretastatin A-4 and hyperthermia. *Radiother Oncol*, 61: 313-320, 2001.
  48. Koukourakis, M. I., Giatromanolaki, A., Sivridis, E., Simopoulos, K., Pastorek, J., Wykoff, C. C., Gatter, K. C., and Harris, A. L. Hypoxia-regulated carbonic anhydrase-9 (CA9) relates to poor vascularization and resistance of squamous cell head and neck cancer to chemoradiotherapy. *Clin Cancer Res*, 7: 3399-4403, 2001.
  49. Lal, A., Peters, H., St Croix, B., Haroon, Z. A., Dewhirst, M. W., Strausberg, R. L., Kaanders, J. H., van der Kogel, A. J., and Riggins, G. J. Transcriptional response to hypoxia in human tumors. *J Natl Cancer Inst*, 93: 1337-1343, 2001.
  50. Lee, Y. M., Jeong, C. H., Koo, S. Y., Son, M. J., Song, H. S., Bae, S. K., Raleigh, J. A., Chung, H. Y., Yoo, M. A., and Kim, K. W. Determination of hypoxic region by hypoxia marker in developing mouse embryos in vivo: a possible signal for vessel development. *Dev. Dyn.*, 220: 175-186, 2001.
  51. Nordmark, M., Loncaster, J., Chou, S. C., Havsteen, H., Lindegaard, J. C., Davidson, S. E., Varia, M., West, C., Hunter, R., Overgaard, J., and Raleigh, J. A. Invasive oxygen measurements and pimonidazole labeling in human cervix carcinoma. *Int J Radiat Oncol Biol Phys*, 49: 581-586, 2001.
  52. Olive, P. L., Aquino-Parsons, C., MacPhail, S. H., Liao, S. Y., Raleigh, J. A., Lerman, M. I., and Stanbridge, E. J. Carbonic anhydrase 9 as an endogenous marker for hypoxic cells in cervical cancer. *Cancer Res*, 61: 8924-8929, 2001.
  53. Olive, P. L., Banath, J. P., and Aquino-Parsons, C. Measuring hypoxia in solid tumours--is there a gold standard? *Acta Oncol*, 40: 917-923, 2001.
  54. Pogue, B. W., Paulsen, K. D., O'Hara, J. A., Wilmot, C. M., and Swartz, H. M. Estimation of oxygen distribution in RIF-1 tumors by diffusion model-based

- interpretation of pimonidazole hypoxia and eppendorf measurements. *Radiat Res*, 155: 15-25, 2001.
55. Raleigh, J. A., Chou, S. C., Bono, E. L., Thrall, D. E., and Varia, M. A. Semiquantitative immunohistochemical analysis for hypoxia in human tumors. *Int J Radiat Oncol Biol Phys*, 49: 569-574, 2001.
  56. Raleigh, J. A., Thrall, D. E., and Varia, M. A. Development and clinical application of pimonidazole as a marker for tumor hypoxia. *Recent Res. Devel. Cancer*, 3: 189-201, 2001.
  57. Schemmer, P., Enomoto, N., Bradford, B. U., Bunzendahl, H., Raleigh, J. A., Lemasters, J. J., and Thurman, R. G. Activated Kupffer cells cause a hypermetabolic state after gentle in situ manipulation of liver in rats. *Am J Physiol Gastrointest Liver Physiol*, 280: G1076-1082, 2001.
  58. Schemmer, P., Enomoto, N., Bradford, B. U., Bunzendahl, H., Raleigh, J. A., and Thurman, R. G. Autonomic nervous system and gut-derived endotoxin: involvement in activation of kupffer cells after in situ organ manipulation. *World J Surg*, 25: 399-406, 2001.
  59. Shabsigh, A., Ghafar, M. A., de la Taille, A., Burchardt, M., Kaplan, S. A., Anastasiadis, A. G., and Buttyan, R. Biomarker analysis demonstrates a hypoxic environment in the castrated rat ventral prostate gland. *J Cell Biochem*, 81: 437-444, 2001.
  60. Suga, S. I., Phillips, M. I., Ray, P. E., Raleigh, J. A., Vio, C. P., Kim, Y. G., Mazzali, M., Gordon, K. L., Hughes, J., and Johnson, R. J. Hypokalemia induces renal injury and alterations in vasoactive mediators that favor salt sensitivity. *Am. J. Physiol. Renal. Physiol.*, 281: F620-629, 2001.
  61. Williams, K. J., Cowen, R. L., and Stratford, I. J. Hypoxia and oxidative stress. Tumour hypoxia--therapeutic considerations. *Breast Cancer Res*, 3: 328-331, 2001.
  62. Zhong, Z., Li, X., Yamashina, S., von Frankenberg, M., Enomoto, N., Ikejima, K., Kolinsky, M., Raleigh, J. A., and Thurman, R. G. Cyclosporin A causes a hypermetabolic state and hypoxia in the liver: prevention by dietary glycine. *J Pharmacol Exp Ther*, 299: 858-865, 2001.
  63. Zips, D., Eicheler, W., Bruchner, K., Jackisch, T., Geyer, P., Petersen, C., van der Kogel, A. J., and Baumann, M. Impact of the tumour bed effect on microenvironment, radiobiological hypoxia and the outcome of fractionated radiotherapy of human FaDu squamous-cell carcinoma growing in the nude mouse. *Int J Radiat Biol*, 77: 1185-1193, 2001.
  64. Arcasoy, M. O., Amin, K., Karayal, A. F., Chou, S. C., Raleigh, J. A., Varia, M. A., and Haroon, Z. A. Functional significance of erythropoietin receptor expression in breast cancer. *Lab Invest*, 82: 911-918, 2002.
  65. Bardag-Gorce, F., French, B. A., Li, J., Riley, N. E., Yuan, Q. X., Valinluck, V., Fu, P., Ingelman-Sundberg, M., Yoon, S., and French, S. W. The importance of cycling of blood alcohol levels in the pathogenesis of experimental alcoholic liver disease in rats. *Gastroenterology*, 123: 325-335, 2002.
  66. Bennewith, K. L., Raleigh, J. A., and Durand, R. E. Orally administered pimonidazole to label hypoxic tumor cells. *Cancer Res*, 62: 6827-6830, 2002.

67. Corpechot, C., Barbu, V., Wendum, D., Chignard, N., Housset, C., Poupon, R., and Rosmorduc, O. Hepatocyte growth factor and c-Met inhibition by hepatic cell hypoxia: a potential mechanism for liver regeneration failure in experimental cirrhosis. *Am J Pathol*, 160: 613-620, 2002.
68. Ghafar, M. A., Anastasiadis, A. G., Olsson, L. E., Chichester, P., Kaplan, S. A., Buttyan, R., and Levin, R. M. Hypoxia and an angiogenic response in the partially obstructed rat bladder. *Lab Invest*, 82: 903-909, 2002.
69. Ghafar, M. A., Shabsigh, A., Chichester, P., Anastasiadis, A. G., Borow, A., Levin, R. M., and Buttyan, R. Effects of chronic partial outlet obstruction on blood flow and oxygenation of the rat bladder. *J Urol*, 167: 1508-1512, 2002.
70. Hale, L. P., Braun, R. D., Gwinn, W. M., Greer, P. K., and Dewhirst, M. W. Hypoxia in the thymus: role of oxygen tension in thymocyte survival. *Am J Physiol Heart Circ Physiol*, 282: H1467-1477, 2002.
71. Harada, H., Hiraoka, M., and Kizaka-Kondoh, S. Antitumor effect of TAT-oxygen-dependent degradation-caspase-3 fusion protein specifically stabilized and activated in hypoxic tumor cells. *Cancer Res*, 62: 2013-2018, 2002.
72. Hoebers, F. J., Janssen, H. L., Olmos, A. V., Sprong, D., Nunn, A. D., Balm, A. J., Hoefnagel, C. A., Begg, A. C., Haustermans, K. M., Ljungkvist, A. S., Bussink, J., Rijken, P. F., Kaanders, J. H., van der Kogel, A. J., Denekamp, J., and Bonn, D. Phase 1 study to identify tumour hypoxia in patients with head and neck cancer using technetium-99m BRU 59-21. Vascular architecture, hypoxia, and proliferation in first-generation xenografts of human head-and-neck squamous cell carcinomas. Why do hypoxic cells behave badly? *Eur J Nucl Med Mol Imaging*, 29: 1206-1211, 2002.
73. Janssen, H. L., Haustermans, K. M., Sprong, D., Blommestijn, G., Hofland, I., Hoebers, F. J., Blijweert, E., Raleigh, J. A., Semenza, G. L., Varia, M. A., Balm, A. J., van Velthuysen, M. L., Delaere, P., Sciort, R., and Begg, A. C. HIF-1alpha, pimonidazole, and iododeoxyuridine to estimate hypoxia and perfusion in human head-and-neck tumors. *Int J Radiat Oncol Biol Phys*, 54: 1537-1549, 2002.
74. Kaanders, J. H., Wijffels, K. I., Marres, H. A., Ljungkvist, A. S., Pop, L. A., van den Hoogen, F. J., de Wilde, P. C., Bussink, J., Raleigh, J. A., and van der Kogel, A. J. Pimonidazole binding and tumor vascularity predict for treatment outcome in head and neck cancer. *Cancer Res*, 62: 7066-7074, 2002.
75. Olive, P. L., Banath, J. P., and Durand, R. E. The range of oxygenation in SiHa tumor xenografts. *Radiat Res*, 158: 159-166, 2002.
76. Olive, P. L., Luo, C. M., and Banath, J. P. Local hypoxia is produced at sites of intratumour injection. *Br J Cancer*, 86: 429-435, 2002.
77. Pedley, R. B., El-Emir, E. b., Flynn, A. A., Boxer, G. M., Dearling, J., Raleigh, J. A., Hill, S. A., Stuart, S., Motha, R., and Begent, R. H. Synergy between vascular targeting agents and antibody-directed therapy. *Int. J. Radiat. Oncol. Biol. Phys.*, 54: 1524-1531, 2002.
78. Rijken, P. F., Peters, J. P., and Van der Kogel, A. J. Quantitative analysis of varying profiles of hypoxia in relation to functional vessels in different human glioma xenograft lines. *Radiat Res*, 157: 626-632, 2002.

79. Rofstad, E. K. and Halsor, E. F. Hypoxia-associated spontaneous pulmonary metastasis in human melanoma xenografts: involvement of microvascular hot spots induced in hypoxic foci by interleukin 8. *Br J Cancer*, 86: 301-308, 2002.
80. Rofstad, E. K., Rasmussen, H., Galappathi, K., Mathiesen, B., Nilsen, K., and Graff, B. A. Hypoxia promotes lymph node metastasis in human melanoma xenografts by up-regulating the urokinase-type plasminogen activator receptor. *Cancer Res.*, 62: 1847-1853, 2002.
81. Rofstad, E. K., Tunheim, S. H., Mathiesen, B., Graff, B. A., Halsor, E. F., Nilsen, K., and Galappathi, K. Pulmonary and lymph node metastasis is associated with primary tumor interstitial fluid pressure in human melanoma xenografts. *Cancer Res*, 62: 661-664, 2002.
82. Yin, M., Zhong, Z., Connor, H. D., Bunzendahl, H., Finn, W. F., Rusyn, I., Li, X., Raleigh, J. A., Mason, R. P., and Thurman, R. G. Protective effect of glycine on renal injury induced by ischemia- reperfusion in vivo. *Am. J. Physiol. Renal. Physiol.*, 282: F417-423, 2002.
83. Airley, R. E., Loncaster, J., Raleigh, J. A., Harris, A. L., Davidson, S. E., Hunter, R. D., West, C. M., and Stratford, I. J. GLUT-1 and CAIX as intrinsic markers of hypoxia in carcinoma of the cervix: Relationship to pimonidazole binding. *Int. J. Cancer*, 104: 85-91, 2003.
84. Azuma, Y., Chou, S. C., Lininger, R. A., Murphy, B. J., Varia, M. A., and Raleigh, J. A. Hypoxia and differentiation in squamous cell carcinomas of the uterine cervix: pimonidazole and involucrin. *Clin Cancer Res*, 9: 4944-4452, 2003.
85. Basile, D. P., Donohoe, D. L., Roethe, K., and Mattson, D. L. Chronic renal hypoxia after acute ischemic injury: effects of L-arginine on hypoxia and secondary damage. *Am J Physiol Renal Physiol*, 284: F338-348, 2003.
86. Bassnett, S. and McNulty, R. The effect of elevated intraocular oxygen on organelle degradation in the embryonic chicken lens. *J. Exp. Biol.*, 206: 4353-4361, 2003.
87. Begg, A. C. Is HIF-1alpha a good marker for tumor hypoxia? *Int. J. Radiat. Oncol. Biol. Phys.*, 56: 917-919, 2003.
88. Blouw, B., Song, H., Tihan, T., Bosze, J., Ferrara, N., Gerber, H. P., Johnson, R. S., and Bergers, G. The hypoxic response of tumors is dependent on their microenvironment. *Cancer Cell*, 4: 133-146, 2003.
89. Bussink, J., Kaanders, J. H., and van der Kogel, A. J. Tumor hypoxia at the micro-regional level: clinical relevance and predictive value of exogenous and endogenous hypoxic cell markers. *Radiother Oncol*, 67: 3-15, 2003.
90. Cachat, F., Lange-Sperandio, B., Chang, A. Y., Kiley, S. C., Thornhill, B. A., Forbes, M. S., and Chevalier, R. L. Ureteral obstruction in neonatal mice elicits segment-specific tubular cell responses leading to nephron loss. *Kidney Int*, 63: 564-575, 2003.
91. Danielsson, B. R., Skold, A. C., Johansson, A., Dillner, B., and Blomgren, B. Teratogenicity by the hERG potassium channel blocking drug almokalant: use of hypoxia marker gives evidence for a hypoxia-related mechanism mediated via embryonic arrhythmia. *Toxicol. Appl. Pharmacol.*, 193: 168-176, 2003.

92. Freeburg, P. B., Robert, B., St John, P. L., and Abrahamson, D. R. Podocyte expression of hypoxia-inducible factor (HIF)-1 and HIF-2 during glomerular development. *J Am Soc Nephrol*, 14: 927-938, 2003.
93. Hoskin, P. J., Sibtain, A., Daley, F. M., and Wilson, G. D. GLUT1 and CAIX as intrinsic markers of hypoxia in bladder cancer: relationship with vascularity and proliferation as predictors of outcome of ARCON. *Br J Cancer*, 89: 1290-1297, 2003.
94. Levin, R. M., O'Connor, L. J., Leggett, R. E., Whitbeck, C., and Chichester, P. Focal hypoxia of the obstructed rabbit bladder wall correlates with intermediate decompensation. *Neurourol Urodyn*, 22: 156-163, 2003.
95. McKim, S. E., Uesugi, T., Raleigh, J. A., McClain, C. J., and Arteel, G. E. Chronic intragastric alcohol exposure causes hypoxia and oxidative stress in the rat pancreas. *Arch Biochem Biophys*, 417: 34-43, 2003.
96. Nordmark, M., Loncaster, J., Aquino-Parsons, C., Chou, S. C., Ladekarl, M., Havsteen, H., Lindegaard, J. C., Davidson, S. E., Varia, M., West, C., Hunter, R., Overgaard, J., and Raleigh, J. A. Measurements of hypoxia using pimonidazole and polarographic oxygen-sensitive electrodes in human cervix carcinomas. *Radiother Oncol*, 67: 35-44, 2003.
97. Petersen, C., Eicheler, W., Frommel, A., Krause, M., Balschukat, S., Zips, D., and Baumann, M. Proliferation and micromilieu during fractionated irradiation of human FaDu squamous cell carcinoma in nude mice. *Int J Radiat Biol*, 79: 469-477, 2003.
98. Samoszuk, M. and Corwin, M. A. Mast cell inhibitor cromolyn increases blood clotting and hypoxia in murine breast cancer. *Int J Cancer*, 107: 159-163, 2003.
99. Sminia, P., Acker, H., Eikesdal, H. P., Kaaijk, P., Enger, P., Slotman, B., and Bjerkvig, R. Oxygenation and response to irradiation of organotypic multicellular spheroids of human glioma. *Anticancer Res.*, 23: 1461-1466, 2003.
100. Vordermark, D. and Brown, J. M. Endogenous markers of tumor hypoxia predictors of clinical radiation resistance? *Strahlenther. Onkol.*, 179: 801-811, 2003.
101. Vordermark, D. and Brown, J. M. Evaluation of hypoxia-inducible factor-1alpha (HIF-1alpha) as an intrinsic marker of tumor hypoxia in U87 MG human glioblastoma: in vitro and xenograft studies. *Int. J. Radiat. Oncol. Biol. Phys.*, 56: 1184-1193, 2003.
102. Zhao, D., Ran, S., Constantinescu, A., Hahn, E. W., and Mason, R. P. Tumor oxygen dynamics: correlation of in vivo MRI with histological findings. *Neoplasia*, 5: 308-318, 2003.
103. Zoula, S., Rijken, P. F., Peters, J. P., Farion, R., Van der Sanden, B. P., Van der Kogel, A. J., Decorps, M., and Remy, C. Pimonidazole binding in C6 rat brain glioma: relation with lipid droplet detection. *Br J Cancer*, 88: 1439-1444, 2003.
104. Bennewith, K. L. and Durand, R. E. Quantifying transient hypoxia in human tumor xenografts by flow cytometry. *Cancer Res*, 64: 6183-6189, 2004.
105. Bhattacharya, A., Toth, K., Mazurchuk, R., Spornyak, J. A., Slocum, H. K., Pendyala, L., Azrak, R., Cao, S., Durrani, F. A., and Rustum, Y. M. Lack of microvessels in well-differentiated regions of human head and neck squamous cell carcinoma A253 associated with functional magnetic resonance imaging

- detectable hypoxia, limited drug delivery, and resistance to irinotecan therapy. *Clin Cancer Res*, *10*: 8005-8017, 2004.
106. Buchler, P., Reber, H. A., Lavey, R. S., Tomlinson, J., Buchler, M. W., Friess, H., and Hines, O. J. Tumor hypoxia correlates with metastatic tumor growth of pancreatic cancer in an orthotopic murine model. *J Surg Res*, *120*: 295-303, 2004.
  107. Cardenas-Navia, L. I., Yu, D., Braun, R. D., Brizel, D. M., Secomb, T. W., and Dewhirst, M. W. Tumor-dependent kinetics of partial pressure of oxygen fluctuations during air and oxygen breathing. *Cancer Res*, *64*: 6010-6017, 2004.
  110. Chou, S. C., Azuma, Y., Varia, M. A., and Raleigh, J. A. Evidence that involucrin, a marker for differentiation, is oxygen regulated in human squamous cell carcinomas. *Br. J. Cancer*, *90*: 728-735, 2004.
  111. Copple, B. L., Rondelli, C. M., Maddox, J. F., Hoglen, N. C., Ganey, P. E., and Roth, R. A. Modes of cell death in rat liver after monocrotaline exposure. *Toxicol Sci*, *77*: 172-182, 2004.
  110. Dearling, J. L., Flynn, A. A., Sutcliffe-Goulden, J., Petrie, I. A., Boden, R., Green, A. J., Boxer, G. M., Begent, R. H., and Pedley, R. B. Analysis of the regional uptake of radiolabeled deoxyglucose analogs in human tumor xenografts. *J. Nucl. Med.*, *45*: 101-107, 2004.
  111. Dubois, L., Landuyt, W., Haustermans, K., Dupont, P., Bormans, G., Vermaelen, P., Flamen, P., Verbeken, E., and Mortelmans, L. Evaluation of hypoxia in an experimental rat tumour model by [(18)F]fluoromisonidazole PET and immunohistochemistry. *Br J Cancer*, *91*: 1947-1954, 2004.
  112. Freeburg, P. B. and Abrahamson, D. R. Divergent expression patterns for hypoxia-inducible factor-1beta and aryl hydrocarbon receptor nuclear transporter-2 in developing kidney. *J Am Soc Nephrol*, *15*: 2569-2578, 2004.
  113. Hoskin, P. J., Sibtain, A., Daley, F. M., Saunders, M. I., and Wilson, G. D. The immunohistochemical assessment of hypoxia, vascularity and proliferation in bladder carcinoma. *Radiother Oncol*, *72*: 159-168, 2004.
  114. Huang, J., Frischer, J. S., New, T., Kim, E. S., Serur, A., Lee, A., Kadenhe-Chiwishe, A., Pollyea, D. A., Yokoi, A., Holash, J., Yancopoulos, G. D., Kandel, J. J., and Yamashiro, D. J. TNP-470 promotes initial vascular sprouting in xenograft tumors. *Mol. Cancer Ther.* *3*: 335-343, 2004.
  115. Hutchison, G. J., Valentine, H. R., Loncaster, J. A., Davidson, S. E., Hunter, R. D., Roberts, S. A., Harris, A. L., Stratford, I. J., Price, P. M., and West, C. M. Hypoxia-inducible factor 1alpha expression as an intrinsic marker of hypoxia: correlation with tumor oxygen, pimonidazole measurements, and outcome in locally advanced carcinoma of the cervix. *Clin Cancer Res*, *10*: 8405-8412, 2004.
  116. Huxham, L. A., Kyle, A. H., Baker, J. H., Nykilchuk, L. K., and Minchinton, A. I. Microregional effects of gemcitabine in HCT-116 xenografts. *Cancer Res*, *64*: 6537-6541, 2004.
  117. Janssen, H. L., Hoebbers, F. J., Sprong, D., Goethals, L., Williams, K. J., Stratford, I. J., Haustermans, K. M., Balm, A. J., and Begg, A. C. Differentiation-associated staining with anti-pimonidazole antibodies in head and neck tumors. *Radiother. Oncol.*, *70*: 91-97, 2004.
  118. Jeong, W. I., Do, S. H., Yun, H. S., Song, B. J., Kim, S. J., Kwak, W. J., Yoo, S. E., Park, H. Y., and Jeong, K. S. Hypoxia potentiates transforming growth factor-

- beta expression of hepatocyte during the cirrhotic condition in rat liver. *Liver Int*, 24: 658-668, 2004.
119. Kostourou, V., Troy, H., Murray, J. F., Cullis, E. R., Whitley, G. S., Griffiths, J. R., and Robinson, S. P. Overexpression of dimethylarginine dimethylaminohydrolase enhances tumor hypoxia: an insight into the relationship of hypoxia and angiogenesis in vivo. *Neoplasia*, 6: 401-411, 2004.
  120. Levin, R., Chichester, P., Levin, S., and Buttyan, R. Role of angiogenesis in bladder response to partial outlet obstruction. *Scand J Urol Nephrol Suppl*: 37-47, 2004.
  121. Manotham, K., Tanaka, T., Matsumoto, M., Ohse, T., Miyata, T., Inagi, R., Kurokawa, K., Fujita, T., and Nangaku, M. Evidence of tubular hypoxia in the early phase in the remnant kidney model. *J. Am. Soc. Nephrol.*, 15: 1277-1288, 2004.
  122. Matsumoto, M., Tanaka, T., Yamamoto, T., Noiri, E., Miyata, T., Inagi, R., Fujita, T., and Nangaku, M. Hypoperfusion of Peritubular Capillaries Induces Chronic Hypoxia before Progression of Tubulointerstitial Injury in a Progressive Model of Rat Glomerulonephritis. *J. Am. Soc. Nephrol.*, 15: 1574-1581, 2004.
  123. Peters, C. L., Morris, C. J., Mapp, P. I., Blake, D. R., Lewis, C. E., and Winrow, V. R. The transcription factors hypoxia-inducible factor 1alpha and Ets-1 colocalize in the hypoxic synovium of inflamed joints in adjuvant-induced arthritis. *Arthritis Rheum.*, 50: 291-296, 2004.
  124. Prabhakaran, K., Sampson, D. A., and Hoehner, J. C. Neuroblastoma survival and death: an in vitro model of hypoxia and metabolic stress. *J. Surg. Res.*, 116: 288-296, 2004.
  125. Rofstad, E. K., Mathiesen, B., and Galappathi, K. Increased metastatic dissemination in human melanoma xenografts after subcurative radiation treatment: radiation-induced increase in fraction of hypoxic cells and hypoxia-induced up-regulation of urokinase-type plasminogen activator receptor. *Cancer Res.*, 64: 13-18, 2004.
  126. Samoszuk, M. K., Walter, J., and Mechetner, E. Improved immunohistochemical method for detecting hypoxia gradients in mouse tissues and tumors. *J Histochem Cytochem*, 52: 837-839, 2004.
  127. Tang, N., Wang, L., Esko, J., Giordano, F. J., Huang, Y., Gerber, H. P., Ferrara, N., and Johnson, R. S. Loss of HIF-1alpha in endothelial cells disrupts a hypoxia-driven VEGF autocrine loop necessary for tumorigenesis. *Cancer Cell*, 6: 485-495, 2004.
  128. van Laarhoven, H. W., Bussink, J., Lok, J., Punt, C. J., Heerschap, A., and van Der Kogel, A. J. Effects of nicotinamide and carbogen in different murine colon carcinomas: immunohistochemical analysis of vascular architecture and microenvironmental parameters. *Int J Radiat Oncol Biol Phys*, 60: 310-321, 2004.
  129. van Wijngaarden, J., de Rooij, K., van Beek, E., Bernsen, H., Que, I., van Hinsbergh, V. W., and Lowik, C. Identification of differentially expressed genes in a renal cell carcinoma tumor model after endostatin-treatment. *Lab Invest*, 84: 1472-1483, 2004.
  130. Yang, L., DeBusk, L. M., Fukuda, K., Fingleton, B., Green-Jarvis, B., Shyr, Y., Matrisian, L. M., Carbone, D. P., and Lin, P. C. Expansion of myeloid immune

- suppressor Gr<sup>+</sup>CD11b<sup>+</sup> cells in tumor-bearing host directly promotes tumor angiogenesis. *Cancer Cell*, 6: 409-421, 2004.
131. Zips, D., Adam, M., Flentje, M., Haase, A., Molls, M., Mueller-Klieser, W., Petersen, C., Philbrook, C., Schmitt, P., Thews, O., Walenta, S., and Baumann, M. Impact of Hypoxia and the Metabolic Microenvironment on Radiotherapy of Solid Tumors Introduction of a Multiinstitutional Research Project. *Strahlenther Onkol*, 180: 609-615, 2004.
  132. Airley, R. E., Phillips, R. M., Evans, A. E., Double, J., Burger, A. M., Feibig, H. H., West, C. M., and Stratford, I. J. Hypoxia-regulated glucose transporter Glut-1 may influence chemosensitivity to some alkylating agents: Results of EORTC (First Translational Award) study of the relevance of tumour hypoxia to the outcome of chemotherapy in human tumour-derived xenografts. *Int J Oncol*, 26: 1477-1484, 2005.
  133. Arcasoy, M. O., Amin, K., Chou, S. C., Haroon, Z. A., Varia, M., and Raleigh, J. A. Erythropoietin and erythropoietin receptor expression in head and neck cancer: relationship to tumor hypoxia. *Clin Cancer Res*, 11: 20-27, 2005.
  134. Asosingh, K., De Raeve, H., de Ridder, M., Storme, G. A., Willems, A., Van Riet, I., Van Camp, B., and Vanderkerken, K. Role of the hypoxic bone marrow microenvironment in 5T2MM murine myeloma tumor progression. *Haematologica*, 90: 810-817, 2005.
  135. Cao, Y., Li, C. Y., Moeller, B. J., Yu, D., Zhao, Y., Dreher, M. R., Shan, S., and Dewhirst, M. W. Observation of incipient tumor angiogenesis that is independent of hypoxia and hypoxia inducible factor-1 activation. *Cancer Res*, 65: 5498-5505, 2005.
  136. Covington, M. D., Bayless, K. J., Burghardt, R. C., Davis, G. E., and Parrish, A. R. Ischemia-induced cleavage of cadherins in NRK cells: evidence for a role of metalloproteinases. *Am J Physiol Renal Physiol*, 289: F280-288, 2005.
  137. Damaser, M. S., Whitbeck, C., Chichester, P., and Levin, R. M. Effect of vaginal distension on blood flow and hypoxia of urogenital organs of the female rat. *J Appl Physiol*, 98: 1884-1890, 2005.
  138. Danielsson, B. R., Johansson, A., Danielsson, C., Azarbayjani, F., Blomgren, B., and Skold, A. C. Phenytoin teratogenicity: Hypoxia marker and effects on embryonic heart rhythm suggest an hERG-related mechanism. *Birth Defects Res A Clin Mol Teratol*, 73: 146-153, 2005.
  139. Doege, K., Heine, S., Jensen, I., Jelkmann, W., and Metzen, E. Inhibition of mitochondrial respiration elevates oxygen concentration but leaves regulation of hypoxia-inducible factor (HIF) intact. *Blood*, 106: 2311-2317, 2005.
  140. Forder, J. P., Munzenmaier, D. H., and Greene, A. S. Angiogenic protection from focal ischemia with angiotensin II type 1 receptor blockade in the rat. *Am J Physiol Heart Circ Physiol*, 288: H1989-1996, 2005.
  141. Gardiner, T. A., Gibson, D. S., de Gooyer, T. E., de la Cruz, V. F., McDonald, D. M., and Stitt, A. W. Inhibition of Tumor Necrosis Factor- $\alpha$  Improves Physiological Angiogenesis and Reduces Pathological Neovascularization in Ischemic Retinopathy. *Am. J. Pathol.*, 166: 637-644, 2005.

142. Harada, H., Kizaka-Kondoh, S., and Hiraoka, M. Optical imaging of tumor hypoxia and evaluation of efficacy of a hypoxia-targeting drug in living animals. *Mol Imaging*, 4: 182-193, 2005.
143. Hofer, S. O., Mitchell, G. M., Penington, A. J., Morrison, W. A., Romeomeeuw, R., Keramidaris, E., Palmer, J., and Knight, K. R. The use of pimonidazole to characterise hypoxia in the internal environment of an in vivo tissue engineering chamber. *Br J Plast Surg*, 58: 1104-1114, 2005.
144. Hoogsteen, I. J., Peeters, W. J., Marres, H. A., Rijken, P. F., van den Hoogen, F. J., van der Kogel, A. J., and Kaanders, J. H. Erythropoietin receptor is not a surrogate marker for tumor hypoxia and does not correlate with survival in head and neck squamous cell carcinomas. *Radiother Oncol*, 76: 213-218, 2005.
145. Janssen, H. L., Haustermans, K. M., Balm, A. J., and Begg, A. C. Hypoxia in head and neck cancer: how much, how important? *Head Neck*, 27: 622-638, 2005.
146. Lewis, C. and Murdoch, C. Macrophage responses to hypoxia: implications for tumor progression and anti-cancer therapies. *Am J Pathol*, 167: 627-635, 2005.
147. Lin, A. D., Mannikarottu, A., Chaudhry, A., Whitbeck, C., Kogan, B. A., Chichester, P., and Levin, R. M. Protective effects of grape suspension on in vivo ischaemia/reperfusion of the rabbit bladder. *BJU Int*, 96: 1397-1402, 2005.
148. Liu, J., Qu, R., Ogura, M., Shibata, T., Harada, H., and Hiraoka, M. Real-time Imaging of Hypoxia-inducible Factor-1 Activity in Tumor Xenografts. *J Radiat Res (Tokyo)*, 46: 93-102, 2005.
149. Ljungkvist, A. S., Bussink, J., Kaanders, J. H., Rijken, P. F., Begg, A. C., Raleigh, J. A., and van der Kogel, A. J. Hypoxic cell turnover in different solid tumor lines. *Int J Radiat Oncol Biol Phys*, 62: 1157-1168, 2005.
150. Lunt, S. J., Telfer, B. A., Fitzmaurice, R. J., Stratford, I. J., and Williams, K. J. Tirapazamine administered as a neoadjuvant to radiotherapy reduces metastatic dissemination. *Clin Cancer Res*, 11: 4212-4216, 2005.
151. Mayer, A., Hockel, M., and Vaupel, P. Carbonic anhydrase IX expression and tumor oxygenation status do not correlate at the microregional level in locally advanced cancers of the uterine cervix. *Clin Cancer Res*, 11: 7220-7225, 2005.
152. Mizukami, Y., Jo, W. S., Duerr, E. M., Gala, M., Li, J., Zhang, X., Zimmer, M. A., Iliopoulos, O., Zukerberg, L. R., Kohgo, Y., Lynch, M. P., Rueda, B. R., and Chung, D. C. Induction of interleukin-8 preserves the angiogenic response in HIF-1alpha-deficient colon cancer cells. *Nat Med*, 11: 992-997, 2005.
153. Murdoch, C. and Lewis, C. E. Macrophage migration and gene expression in response to tumor hypoxia. *Int J Cancer*, 117: 701-708, 2005.
154. Nelson, D. W., Cao, H., Zhu, Y., Sunar-Reeder, B., Choi, C. Y., Faix, J. D., Brown, J. M., Koong, A. C., Giaccia, A. J., and Le, Q. T. A noninvasive approach for assessing tumor hypoxia in xenografts: developing a urinary marker for hypoxia. *Cancer Res*, 65: 6151-6158, 2005.
155. O'Donoghue J, A., Zanzonico, P., Pugachev, A., Wen, B., Smith-Jones, P., Cai, S., Burnazi, E., Finn, R. D., Burgman, P., Ruan, S., Lewis, J. S., Welch, M. J., Ling, C. C., and Humm, J. L. Assessment of regional tumor hypoxia using (18)F-fluoromisonidazole and (64)Cu(II)-diacetyl-bis(N4-methylthiosemicarbazone) positron emission tomography: Comparative study featuring microPET imaging, Po(2) probe measurement, autoradiography, and fluorescent microscopy in the

- R3327-AT and FaDu rat tumor models. *Int J Radiat Oncol Biol Phys*, *61*: 1493-1502, 2005.
156. Pugachev, A., Ruan, S., Carlin, S., Larson, S. M., Campa, J., Ling, C. C., and Humm, J. L. Dependence of FDG uptake on tumor microenvironment. *Int J Radiat Oncol Biol Phys*, *62*: 545-553, 2005.
  157. Rofstad, E. K., Mathiesen, B., Henriksen, K., Kindem, K., and Galappathi, K. The Tumor Bed Effect: increased metastatic dissemination from hypoxia-Induced up-regulation of metastasis-promoting gene products. *Cancer Res*, *65*: 2387-2396, 2005.
  158. Rosenberger, C., Heyman, S. N., Rosen, S., Shina, A., Goldfarb, M., Griethe, W., Frei, U., Reinke, P., Bachmann, S., and Eckardt, K. U. Up-regulation of HIF in experimental acute renal failure: Evidence for a protective transcriptional response to hypoxia. *Kidney Int*, *67*: 531-542, 2005.
  159. Schuurin, J., Bussink, J., Bernsen, H. J., Peeters, W., and van Der Kogel, A. J. Irradiation combined with SU5416: microvascular changes and growth delay in a human xenograft glioblastoma tumor line. *Int J Radiat Oncol Biol Phys*, *61*: 529-534, 2005.
  160. Sobhanifar, S., Aquino-Parsons, C., Stanbridge, E. J., and Olive, P. Reduced Expression of Hypoxia-Inducible Factor-1 {alpha} in Perinecrotic Regions of Solid Tumors. *Cancer Res*, *65*: 7259-7266, 2005.
  161. Solomon, B., Binns, D., Roselt, P., Weibe, L. I., McArthur, G. A., Cullinane, C., and Hicks, R. J. Modulation of intratumoral hypoxia by the epidermal growth factor receptor inhibitor gefitinib detected using small animal PET imaging. *Mol Cancer Ther*, *4*: 1417-1422, 2005.
  162. Thorwarth, D., Eschmann, S. M., Paulsen, F., and Alber, M. A kinetic model for dynamic [(18)F]-Fmiso PET data to analyse tumour hypoxia. *Phys Med Biol*, *50*: 2209-2224, 2005.
  163. Troost, E. G., Bussink, J., Kaanders, J. H., van Eerd, J., Peters, J. P., Rijken, P. F., Boerman, O. C., and van der Kogel, A. J. Comparison of different methods of CAIX quantification in relation to hypoxia in three human head and neck tumor lines. *Radiother Oncol*, *76*: 194-199, 2005.
  164. van der Schaft, D. W. J., Hillen, F., Pauwels, P., Kirschmann, D. A., Castermans, K., oude Egbrink, M. G. A., Tran, M. G. B., Sciot, R., Hauben, E., Hogendoorn, P. C. W., Delattre, O., Maxwell, P. H., Hendrix, M. J. C., and Griffioen, A. W. Tumor Cell Plasticity in Ewing Sarcoma, an Alternative Circulatory System Stimulated by Hypoxia. *Cancer Res*, *65*: 11520-11528, 2005.
  165. van Herpen, C. M., Bussink, J., van der Kogel, A. J., Peeters, W. J., van der Voort, R., van Schijndel, A., de Wilde, P. C., Adema, G. J., and de Mulder, P. H. Interleukin-12 has no effect on vascular density, perfusion, hypoxia and proliferation of an implanted human squamous cell carcinoma xenograft tumour despite up-regulation of ICAM-1. *Anticancer Res*, *25*: 1015-10121, 2005.
  166. van Laarhoven, H. W., de Geus-Oei, L. F., Wiering, B., Lok, J., Rijpkema, M., Kaanders, J. H., Krabbe, P. F., Ruers, T., Punt, C. J., van der Kogel, A. J., Oyen, W. J., and Heerschap, A. Gadopentetate dimeglumine and FDG uptake in liver metastases of colorectal carcinoma as determined with MR imaging and PET. *Radiology*, *237*: 181-188, 2005.

167. Villanueva, S., Cespedes, C., and Vio, C. P. Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. *Am J Physiol Regul Integr Comp Physiol*, 2005.
168. Vordermark, D., Kraft, P., Katzer, A., Bolling, T., Willner, J., and Flentje, M. Glucose requirement for hypoxic accumulation of hypoxia-inducible factor-1alpha (HIF-1alpha). *Cancer Lett*, 230: 122-133, 2005.
169. Wang, P. X. and Sanders, P. W. Mechanism of hypertensive nephropathy in the Dahl/Rapp rat: a primary disorder of vascular smooth muscle. *Am J Physiol Renal Physiol*, 288: F236-242, 2005.
170. Yahiro, T., Masui, S., Kubota, N., Yamada, K., Kobayashi, A., and Kishii, K. Effects of hypoxic cell radiosensitizer doranidazole (PR-350) on the radioresponse of murine and human tumor cells in vitro and in vivo. *J Radiat Res (Tokyo)*, 46: 363-372, 2005.
171. Yaromina, A., Holscher, T., Eicheler, W., Rosner, A., Krause, M., Hessel, F., Petersen, C., Thames, H. D., Baumann, M., and Zips, D. Does heterogeneity of pimonidazole labelling correspond to the heterogeneity of radiation-response of FaDu human squamous cell carcinoma? *Radiother Oncol*, 76: 206-212, 2005.
172. Badger, W. J., Whitbeck, C., Kogan, B., Chichester, P., and Levin, R. M. The immediate effect of castration on female rabbit bladder blood flow and tissue oxygenation. *Urol Int*, 76: 264-268, 2006.
173. Carnell, D. M., Smith, R. E., Daley, F. M., Saunders, M. I., Bentzen, S. M., and Hoskin, P. J. An immunohistochemical assessment of hypoxia in prostate carcinoma using pimonidazole: Implications for radioresistance. *Int J Radiat Oncol Biol Phys*, 65: 91-99, 2006.
174. Ceelen, W., Smeets, P., Backes, W., Van Damme, N., Boterberg, T., Demetter, P., Bouckenoghe, I., De Visschere, M., Peeters, M., and Pattyn, P. Noninvasive monitoring of radiotherapy-induced microvascular changes using dynamic contrast enhanced magnetic resonance imaging (DCE-MRI) in a colorectal tumor model. *Int J Radiat Oncol Biol Phys*, 64: 1188-1196, 2006.
175. Cheema, A. N., Hong, T., Nili, N., Segev, A., Moffat, J. G., Lipson, K. E., Howlett, A. R., Holdsworth, D. W., Cole, M. J., Qiang, B., Kolodgie, F., Virmani, R., Stewart, D. J., and Strauss, B. H. Adventitial microvessel formation after coronary stenting and the effects of SU11218, a tyrosine kinase inhibitor. *J Am Coll Cardiol*, 47: 1067-1075, 2006.
176. Danylkova, N. O., Pomeranz, H. D., Alcala, S. R., and McLoon, L. K. Histological and morphometric evaluation of transient retinal and optic nerve ischemia in rat. *Brain Res*, 1096: 20-29, 2006.
177. Dorow, D. S., Cullinane, C., Conus, N., Roselt, P., Binns, D., McCarthy, T. J., McArthur, G. A., and Hicks, R. J. Multi-tracer small animal PET imaging of the tumour response to the novel pan-Erb-B inhibitor CI-1033. *Eur J Nucl Med Mol Imaging*: 1-12, 2006.
178. Franco, M., Man, S., Chen, L., Emmenegger, U., Shaked, Y., Cheung, A. M., Brown, A. S., Hicklin, D. J., Foster, F. S., and Kerbel, R. S. Targeted anti-vascular endothelial growth factor receptor-2 therapy leads to short-term and long-term impairment of vascular function and increase in tumor hypoxia. *Cancer Res*, 66: 3639-3648, 2006.

179. Goethals, L., Debucquoy, A., Perneel, C., Geboes, K., Ectors, N., De Schutter, H., Penninckx, F., McBride, W. H., Begg, A. C., and Haustermans, K. M. Hypoxia in human colorectal adenocarcinoma: Comparison between extrinsic and potential intrinsic hypoxia markers. *Int J Radiat Oncol Biol Phys*, *65*: 246-254, 2006.
180. Goldfarb, M., Rosenberger, C., Abassi, Z., Shina, A., Zilbersat, F., Eckardt, K. U., Rosen, S., and Heyman, S. N. Acute-on-chronic renal failure in the rat: functional compensation and hypoxia tolerance. *Am J Nephrol*, *26*: 22-33, 2006.
181. Jankovic, B., Aquino-Parsons, C., Raleigh, J. A., Stanbridge, E. J., Durand, R. E., Banath, J. P., MacPhail, S. H., and Olive, P. L. Comparison between pimonidazole binding, oxygen electrode measurements, and expression of endogenous hypoxia markers in cancer of the uterine cervix. *Cytometry B Clin Cytom*, *70*: 45-55, 2006.
182. Kabuubi, P., Loncaster, J. A., Davidson, S. E., Hunter, R. D., Kobylecki, C., Stratford, I. J., and West, C. M. No relationship between thymidine phosphorylase (TP, PD-ECGF) expression and hypoxia in carcinoma of the cervix. *Br J Cancer*, *94*: 115-120, 2006.
183. Kirkpatrick, J. P., Hardee, M. E., Snyder, S. A., Peltz, C. M., Zhao, Y., Brizel, D. M., Dewhirst, M. W., and Blackwell, K. L. The effect of darbepoetin alfa on growth, oxygenation and radioresponsiveness of a breast adenocarcinoma. *Radiat Res*, *165*: 192-201, 2006.
184. Kleiter, M. M., Thrall, D. E., Malarkey, D. E., Ji, X., Lee, D. Y., Chou, S. C., and Raleigh, J. A. A comparison of oral and intravenous pimonidazole in canine tumors using intravenous CCI-103F as a control hypoxia marker. *Int J Radiat Oncol Biol Phys*, *64*: 592-602, 2006.
185. Lipnik, K., Greco, O., Scott, S., Knapp, E., Mayrhofer, E., Rosenfellner, D., Gunzburg, W. H., Salmons, B., and Hohenadl, C. Hypoxia- and radiation-inducible, breast cell-specific targeting of retroviral vectors. *Virology*, *349*: 121-133, 2006.
186. Ljungkvist, A. S., Bussink, J., Kaanders, J. H., Wiedenmann, N. E., Vlasman, R., and van der Kogel, A. J. Dynamics of hypoxia, proliferation and apoptosis after irradiation in a murine tumor model. *Radiat Res*, *165*: 326-336, 2006.
187. Morani, A., Barros, R. P., Imamov, O., Hultenby, K., Arner, A., Warner, M., and Gustafsson, J. A. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta<sup>-/-</sup>) mice. *Proc Natl Acad Sci U S A*, *103*: 7165-7169, 2006.
188. Nanka, O., Valasek, P., Dvorakova, M., and Grim, M. Experimental hypoxia and embryonic angiogenesis. *Dev Dyn*, *235*: 723-733, 2006.
189. van Laarhoven, H. W., Kaanders, J. H., Lok, J., Peeters, W. J., Rijken, P. F., Wiering, B., Ruers, T. J., Punt, C. J., Heerschap, A., and van der Kogel, A. J. Hypoxia in relation to vasculature and proliferation in liver metastases in patients with colorectal cancer. *Int J Radiat Oncol Biol Phys*, *64*: 473-482, 2006.
190. Villanueva, S., Cespedes, C., and Vio, C. P. Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. *Am J Physiol Regul Integr Comp Physiol*, *290*: R861-870, 2006.
191. Yuan, H., Schroeder, T., Bowsher, J. E., Hedlund, L. W., Wong, T., and Dewhirst, M. W. Intertumoral differences in hypoxia selectivity of the PET

- imaging agent  $^{64}\text{Cu}(\text{II})$ -diacetyl-bis(N4-methylthiosemicarbazone). *J Nucl Med*, 47: 989-998, 2006.
192. Aly, S., Wagner, K., Keller, C., Malm, S., Malzan, A., Brandau, S., Bange, F. C., and Ehlers, S. Oxygen status of lung granulomas in *Mycobacterium tuberculosis*-infected mice. *J Pathol*, 210: 298-305, 2006.
  193. Copple, B. L., Roth, R. A., and Ganey, P. E. Anticoagulation and inhibition of nitric oxide synthase influence hepatic hypoxia after monocrotaline exposure. *Toxicology*, 225: 128-137, 2006.
  194. de Gooyer, T. E., Stevenson, K. A., Humphries, P., Simpson, D. A., Curtis, T. M., Gardiner, T. A., and Stitt, A. W. Rod photoreceptor loss in Rho<sup>-/-</sup> mice reduces retinal hypoxia and hypoxia-regulated gene expression. *Invest Ophthalmol Vis Sci*, 47: 5553-5560, 2006.
  195. de Gooyer, T. E., Stevenson, K. A., Humphries, P., Simpson, D. A., Gardiner, T. A., and Stitt, A. W. Retinopathy is reduced during experimental diabetes in a mouse model of outer retinal degeneration. *Invest Ophthalmol Vis Sci*, 47: 5561-5568, 2006.
  196. De Schutter, H., Barbe, B., Spaepen, M., Begg, A. C., Balm, A., Gregoire, V., Haustermans, K., Mahy, P., Vander Poorten, V., and Nuyts, S. Microsatellite alterations in head and neck squamous cell carcinoma and relation to expression of pimonidazole, CA IX and GLUT-1. *Radiother Oncol*, 80: 143-150, 2006.
  197. Durand, R. E. and Aquino-Parsons, C. The fate of hypoxic (pimonidazole-labelled) cells in human cervix tumours undergoing chemo-radiotherapy. *Radiother Oncol*, 80(2):138-142, 2006.
  198. Gookin, J. L., Copple, C. N., Papich, M. G., Poore, M. F., Stauffer, S. H., Birkenheuer, A. J., Twedt, D. C., and Levy, M. G. Efficacy of ronidazole for treatment of feline *Tritrichomonas foetus* infection. *J Vet Intern Med*, 20: 536-543, 2006.
  199. Greco, O., Joiner, M. C., Doleh, A., Powell, A. D., Hillman, G. G., and Scott, S. D. Hypoxia- and radiation-activated Cre/loxP 'molecular switch' vectors for gene therapy of cancer. *Gene Ther*, 13: 206-215, 2006.
  200. Huxham, L. A., Kyle, A. H., Baker, J. H., McNicol, K. L., and Minchinton, A. I. Tirapazamine causes vascular dysfunction in HCT-116 tumour xenografts. *Radiother Oncol*, 78: 138-145, 2006.
  201. Jensen, R. L. Hypoxia in the tumorigenesis of gliomas and as a potential target for therapeutic measures. *Neurosurg Focus*, 20: E24, 2006.
  202. Jonathan, R. A., Wijffels, K. I., Peeters, W., de Wilde, P. C., Marres, H. A., Merckx, M. A., Oosterwijk, E., van der Kogel, A. J., and Kaanders, J. H. The prognostic value of endogenous hypoxia-related markers for head and neck squamous cell carcinomas treated with ARCON. *Radiother Oncol*, 79: 288-297, 2006.
  203. Kang, S. H., Cho, H. T., Devi, S., Zhang, Z., Escuin, D., Liang, Z., Mao, H., Brat, D. J., Olson, J. J., Simons, J. W., Lavalley, T. M., Giannakakou, P., Van Meir, E. G., and Shim, H. Antitumor effect of 2-methoxyestradiol in a rat orthotopic brain tumor model. *Cancer Res*, 66: 11991-11997, 2006.
  204. Khan, Z., Michalopoulos, G. K., and Stolz, D. B. Peroxisomal localization of hypoxia-inducible factors and hypoxia-inducible factor regulatory hydroxylases in

- primary rat hepatocytes exposed to hypoxia-reoxygenation. *Am J Pathol*, *169*: 1251-1269, 2006.
205. Laderoute, K. R., Amin, K., Calaoagan, J. M., Knapp, M., Le, T., Orduna, J., Foretz, M., and Viollet, B. 5'-AMP-activated protein kinase (AMPK) is induced by low-oxygen and glucose deprivation conditions found in solid-tumor microenvironments. *Mol Cell Biol*, *26*: 5336-5347, 2006.
  206. Lokmic, Z., Darby, I. A., Thompson, E. W., and Mitchell, G. M. Time course analysis of hypoxia, granulation tissue and blood vessel growth, and remodeling in healing rat cutaneous incisional primary intention wounds. *Wound Repair Regen*, *14*: 277-288, 2006.
  207. Minchinton, A. I. and Tannock, I. F. Drug penetration in solid tumours. *Nat Rev Cancer*, *6*: 583-592, 2006.
  208. Nordmark, M., Loncaster, J., Aquino-Parsons, C., Chou, S. C., Gebiski, V., West, C., Lindegaard, J. C., Havsteen, H., Davidson, S. E., Hunter, R., Raleigh, J. A., and Overgaard, J. The prognostic value of pimonidazole and tumour pO<sub>2</sub> in human cervix carcinomas after radiation therapy: A prospective international multi-center study. *Radiother Oncol*, *80*: 123-131, 2006.
  209. Noto, T., Furuichi, Y., Ishiye, M., Matsuoka, N., Aramori, I., Mutoh, S., Yanagihara, T., and Manabe, N. Temporal and topographic profiles of tissue hypoxia following transient focal cerebral ischemia in rats. *J Vet Med Sci*, *68*: 803-807, 2006.
  210. Papandreou, I., Cairns, R. A., Fontana, L., Lim, A. L., and Denko, N. C. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab*, *3*: 187-197, 2006.
  211. Pugachev, A., Carlin, S., Ling, C., and Humm, J. SU-FF-J-89: Intratumoral Pattern of FDG Uptake in Human Xenograft Models. *Medical Physics* *33*: 2040-, 2006.
  212. Rosenberger, C., Rosen, S., Shina, A., Bernhardt, W., Wiesener, M. S., Frei, U., Eckardt, K. U., and Heyman, S. N. Hypoxia-inducible factors and tubular cell survival in isolated perfused kidneys. *Kidney Int*, *70*: 60-70, 2006.
  213. Schaffer, C. B., Friedman, B., Nishimura, N., Schroeder, L. F., Tsai, P. S., Ebner, F. F., Lyden, P. D., and Kleinfeld, D. Two-photon imaging of cortical surface microvessels reveals a robust redistribution in blood flow after vascular occlusion. *PLoS Biol*, *4*: e22, 2006.
  214. De Schutter, H. D., Barbe, B., Spaepen, M., Begg, A. C., Balm, A., Gregoire, V., Haustermans, K., Mahy, P., Poorten, V. V., and Nuyts, S. Microsatellite alterations in head and neck squamous cell carcinoma and relation to expression of pimonidazole, CA IX and GLUT-1. *Radiother Oncol*, *80*(2):143-150 2006.
  215. Stockwin, L. H., Blonder, J., Bumke, M. A., Lucas, D. A., Chan, K. C., Conrads, T. P., Issaq, H. J., Veenstra, T. D., Newton, D. L., and Rybak, S. M. Proteomic analysis of plasma membrane from hypoxia-adapted malignant melanoma. *J Proteome Res*, *5*: 2996-3007, 2006.
  216. Tanaka, H., Yamamoto, M., Hashimoto, N., Miyakoshi, M., Tamakawa, S., Yoshie, M., Tokusashi, Y., Yokoyama, K., Yaginuma, Y., and Ogawa, K. Hypoxia-independent overexpression of hypoxia-inducible factor 1alpha as an

- early change in mouse hepatocarcinogenesis. *Cancer Res*, *66*: 11263-11270, 2006.
217. Tanaka, T., Kato, H., Kojima, I., Ohse, T., Son, D., Tawakami, T., Yatagawa, T., Inagi, R., Fujita, T., and Nangaku, M. Hypoxia and expression of hypoxia-inducible factor in the aging kidney. *J Gerontol A Biol Sci Med Sci*, *61*: 795-805, 2006.
218. Thorwarth, D., Eschmann, S. M., Holzner, F., Paulsen, F., and Alber, M. Combined uptake of [18F]FDG and [18F]FMISO correlates with radiation therapy outcome in head-and-neck cancer patients. *Radiother Oncol*, *80*: 151-156, 2006.
219. Troost, E. G., Laverman, P., Kaanders, J. H., Philippens, M., Lok, J., Oyen, W. J., van der Kogel, A. J., Boerman, O. C., and Bussink, J. Imaging hypoxia after oxygenation-modification: Comparing [(18)F]FMISO autoradiography with pimonidazole immunohistochemistry in human xenograft tumors. *Radiother Oncol*, 2006.
220. Vignozzi, L., Filippi, S., Morelli, A., Ambrosini, S., Luconi, M., Vannelli, G. B., Donati, S., Crescioli, C., Zhang, X. H., Mirone, V., Forti, G., and Maggi, M. Effect of chronic tadalafil administration on penile hypoxia induced by cavernous neurotomy in the rat. *J Sex Med*, *3*: 419-431, 2006.
221. Webster, W. S., Howe, A. M., Abela, D., and Oakes, D. J. The relationship between cleft lip, maxillary hypoplasia, hypoxia and phenytoin. *Curr Pharm Des*, *12*: 1431-1448, 2006.
222. Yaromina, A., Zips, D., Thames, H. D., Eicheler, W., Krause, M., Rosner, A., Haase, M., Petersen, C., Raleigh, J. A., Quennet, V., Walenta, S., Mueller-Klieser, W., and Baumann, M. Pimonidazole labelling and response to fractionated irradiation of five human squamous cell carcinoma (hSCC) lines in nude mice: The need for a multivariate approach in biomarker studies. *Radiother Oncol*, *81*: 122-129, 2006.
223. Zampino, M., Yuzhakova, M., Hansen, J., McKinney, R. D., Goldspink, P. H., Geenen, D. L., and Buttrick, P. M. Sex-related dimorphic response of HIF-1 alpha expression in myocardial ischemia. *Am J Physiol Heart Circ Physiol*, *291*: H957-964, 2006.
224. Aly, S., Laskay, T., Mages, J., Malzan, A., Lang, R., and Ehlers, S. Interferon-gamma-dependent mechanisms of mycobacteria-induced pulmonary immunopathology: the role of angiostasis and CXCR3-targeted chemokines for granuloma necrosis. *J Pathol*, *212*: 295-305, 2007.
225. Bernhardt, W. M., Wiesener, M. S., Weidemann, A., Schmitt, R., Weichert, W., Lechler, P., Campean, V., Ong, A. C., Willam, C., Gretz, N., and Eckardt, K. U. Involvement of hypoxia-inducible transcription factors in polycystic kidney disease. *Am J Pathol*, *170*: 830-842, 2007.
226. Brown, D. A., MacLellan, W. R., Laks, H., Dunn, J. C., Wu, B. M., and Beygui, R. E. Analysis of oxygen transport in a diffusion-limited model of engineered heart tissue. *Biotechnol Bioeng*, *97*: 962-975, 2007.
227. Burgu, B., Medina Ortiz, W. E., Pitera, J. E., Woolf, A. S., and Wilcox, D. T. Vascular endothelial growth factor mediates hypoxic stimulated embryonic bladder growth in organ culture. *J Urol*, *177*: 1552-1557, 2007.

228. Bussink, J., Kaanders, J. H., and van der Kogel, A. J. Microenvironmental transformations by VEGF- and EGF-receptor inhibition and potential implications for responsiveness to radiotherapy. *Radiother Oncol*, 82: 10-17, 2007.
229. Chang, Y. S., Adnane, J., Trail, P. A., Levy, J., Henderson, A., Xue, D., Bortolon, E., Ichetovkin, M., Chen, C., McNabola, A., Wilkie, D., Carter, C. A., Taylor, I. C., Lynch, M., and Wilhelm, S. Sorafenib (BAY 43-9006) inhibits tumor growth and vascularization and induces tumor apoptosis and hypoxia in RCC xenograft models. *Cancer Chemother Pharmacol*, 59: 561-574, 2007.
230. Cheluvappa, R., Hilmer, S. N., Kwun, S. Y., Cogger, V. C., and DG, L. E. C. Effects of old age on hepatocyte oxygenation. *Ann N Y Acad Sci*, 1114: 88-92, 2007.
231. Cheluvappa, R., Hilmer, S. N., Kwun, S. Y., Jamieson, H. A., O'Reilly, J. N., Muller, M., Cogger, V. C., and Le Couteur, D. G. The effect of old age on liver oxygenation and the hepatic expression of VEGF and VEGFR2. *Exp Gerontol*, 42: 1012-1019, 2007.
232. Degrossoli, A., Bosetto, M. C., Lima, C. B., and Giorgio, S. Expression of hypoxia-inducible factor 1alpha in mononuclear phagocytes infected with *Leishmania amazonensis*. *Immunol Lett*, 114: 119-125, 2007.
233. Deng, X., Luyendyk, J. P., Zou, W., Lu, J., Malle, E., Ganey, P. E., and Roth, R. A. Neutrophil interaction with the hemostatic system contributes to liver injury in rats cotreated with lipopolysaccharide and ranitidine. *J Pharmacol Exp Ther*, 322: 852-861, 2007.
234. Donadiou, E., Hamdi, W., Deveze, A., Lucciano, M., Lavieille, J. P., Magnan, J., and Riva, C. Improved cryosections and specific immunohistochemical methods for detecting hypoxia in mouse and rat cochleae. *Acta Histochem*, 109: 177-184, 2007.
235. Echevarria, M., Munoz-Cabello, A. M., Sanchez-Silva, R., Toledo-Aral, J. J., and Lopez-Barneo, J. Development of cytosolic hypoxia and hypoxia-inducible factor stabilization are facilitated by aquaporin-1 expression. *J Biol Chem*, 282: 30207-30215, 2007.
236. Fang, Y., Sullivan, R., and Graham, C. H. Confluence-dependent resistance to doxorubicin in human MDA-MB-231 breast carcinoma cells requires hypoxia-inducible factor-1 activity. *Exp Cell Res*, 313: 867-877, 2007.
237. Frangoulis, M., Georgiou, P., Chrisostomidis, C., Perrea, D., Dontas, I., Kavantzias, N., Kostakis, A., and Papadopoulos, O. Rat epigastric flap survival and VEGF expression after local copper application. *Plast Reconstr Surg*, 119: 837-843, 2007.
238. Gabellieri, C., Eykyn, T. R., and Leach, M. O. Conformational exchange in pimonidazole--a hypoxia marker. *Magn Reson Chem*, 45: 621-623, 2007.
239. Gonzalez, A. M., Yazici, I., Kusza, K., and Siemionow, M. Effects of fresh versus banked blood transfusions on microcirculatory hemodynamics and tissue oxygenation in the rat cremaster model. *Surgery*, 141: 630-639, 2007.
240. Hoogsteen, I. J., Marres, H. A., Bussink, J., van der Kogel, A. J., and Kaanders, J. H. Tumor microenvironment in head and neck squamous cell carcinomas: Predictive value and clinical relevance of hypoxic markers. A review. *Head Neck*, 2007.

241. Hoskin, P. J., Carnell, D. M., Taylor, N. J., Smith, R. E., Stirling, J. J., Daley, F. M., Saunders, M. I., Bentzen, S. M., Collins, D. J., d'Arcy, J. A., and Padhani, A. P. Hypoxia in prostate cancer: correlation of BOLD-MRI with pimonidazole immunohistochemistry-initial observations. *Int J Radiat Oncol Biol Phys*, *68*: 1065-1071, 2007.
242. Hosogai, N., Fukuhara, A., Oshima, K., Miyata, Y., Tanaka, S., Segawa, K., Furukawa, S., Tochino, Y., Komuro, R., Matsuda, M., and Shimomura, I. Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes*, *56*: 901-911, 2007.
243. Iheagwara, K. N., Thom, S. R., Deutschman, C. S., and Levy, R. J. Myocardial cytochrome oxidase activity is decreased following carbon monoxide exposure. *Biochim Biophys Acta*, 2007.
244. Imamura, R., Moriyama, T., Isaka, Y., Namba, Y., Ichimaru, N., Takahara, S., and Okuyama, A. Erythropoietin protects the kidneys against ischemia reperfusion injury by activating hypoxia inducible factor-1alpha. *Transplantation*, *83*: 1371-1379, 2007.
245. Kaneko, H., Yu, D., and Miura, M. Overexpression of IGF-I receptor in HeLa cells enhances in vivo radioresponse. *Biochem Biophys Res Commun*, *363*: 937-941, 2007.
246. Kim, Y. J., Kang, H. H., Ahn, J. H., and Chung, J. W. Hypoxic changes in the central nervous system of noise-exposed mice. *Acta Otolaryngol Suppl*: 73-77, 2007.
247. Krause, M., Prager, J., Zhou, X., Yaromina, A., Dorfler, A., Eicheler, W., and Baumann, M. EGFR-TK inhibition before radiotherapy reduces tumour volume but does not improve local control: differential response of cancer stem cells and nontumorigenic cells? *Radiother Oncol*, *83*: 316-325, 2007.
248. Le, Q. T. Identifying and targeting hypoxia in head and neck cancer: a brief overview of current approaches. *Int J Radiat Oncol Biol Phys*, *69*: S56-58, 2007.
249. Lenaerts, A. J., Hoff, D., Aly, S., Ehlers, S., Andries, K., Cantarero, L., Orme, I. M., and Basaraba, R. J. Location of persisting mycobacteria in a Guinea pig model of tuberculosis revealed by r207910. *Antimicrob Agents Chemother*, *51*: 3338-3345, 2007.
250. Levesque, J. P., Winkler, I. G., Hendy, J., Williams, B., Helwani, F., Barbier, V., Nowlan, B., and Nilsson, S. K. Hematopoietic Progenitor Cell Mobilization Results in Hypoxia with Increased Hypoxia-Inducible Transcription Factor-1 {alpha} and Vascular Endothelial Growth Factor A in Bone Marrow. *Stem Cells*, *25*: 1954-1965, 2007.
251. Li, X. F., Carlin, S., Urano, M., Russell, J., Ling, C. C., and O'Donoghue, J. A. Visualization of hypoxia in microscopic tumors by immunofluorescent microscopy. *Cancer Res*, *67*: 7646-7653, 2007.
252. Ljungkvist, A. S., Bussink, J., Kaanders, J. H., and van der Kogel, A. J. Dynamics of tumor hypoxia measured with bioreductive hypoxic cell markers. *Radiat Res*, *167*: 127-145, 2007.
253. Matsumoto, K., Szajek, L., Krishna, M. C., Cook, J. A., Seidel, J., Grimes, K., Carson, J., Sowers, A. L., English, S., Green, M. V., Bacharach, S. L., Eckelman, W. C., and Mitchell, J. B. The influence of tumor oxygenation on hypoxia

- imaging in murine squamous cell carcinoma using [64Cu]Cu-ATSM or [18F]Fluoromisonidazole positron emission tomography. *Int J Oncol*, 30: 873-881, 2007.
254. Noto, T., Furuichi, Y., Ishiye, M., Matsuoka, N., Aramori, I., Mutoh, S., and Yanagihara, T. Tacrolimus (FK506) limits accumulation of granulocytes and platelets and protects against brain damage after transient focal cerebral ischemia in rat. *Biol Pharm Bull*, 30: 313-317, 2007.
255. Olmsted-Davis, E., Gannon, F. H., Ozen, M., Ittmann, M. M., Gugala, Z., Hipp, J. A., Moran, K. M., Fouletier-Dilling, C. M., Schumara-Martin, S., Lindsey, R. W., Heggeness, M. H., Brenner, M. K., and Davis, A. R. Hypoxic adipocytes pattern early heterotopic bone formation. *Am J Pathol*, 170: 620-632, 2007.
256. Papadopoulou, M. V., Bloomer, W. D., Taylor, A. P., Hernandez, M., Blumenthal, R. D., and Hollingshead, M. G. Advantage of combining NLCQ-1 (NSC 709257) with radiation in treatment of human head and neck xenografts. *Radiat Res*, 168: 65-71, 2007.
257. Parmar, K., Mauch, P., Vergilio, J. A., Sackstein, R., and Down, J. D. Distribution of hematopoietic stem cells in the bone marrow according to regional hypoxia. *Proc Natl Acad Sci U S A*, 104: 5431-5436, 2007.
258. Plock, J., Frese, S., Keogh, A., Bisch-Knaden, S., Ayuni, E., Corazza, N., Weikert, C., Jakob, S., Erni, D., Dufour, J. F., Brunner, T., Candinas, D., and Stroka, D. Activation of non-ischemic, hypoxia-inducible signalling pathways up-regulate cytoprotective genes in the murine liver. *J Hepatol*, 47: 538-545, 2007.
259. Pringle, K. G., Kind, K. L., Thompson, J. G., and Roberts, C. T. Complex Interactions Between Hypoxia Inducible Factors, Insulin-Like Growth Factor-II and Oxygen in Early Murine Trophoblasts. *Placenta*, 28: 11-12, 2007.
260. Riva, C., Donadieu, E., Magnan, J., and Lavieille, J. P. Age-related hearing loss in CD/1 mice is associated to ROS formation and HIF target proteins up-regulation in the cochlea. *Exp Gerontol*, 42: 327-336, 2007.
261. Rofstad, E. K., Galappathi, K., Mathiesen, B., and Ruud, E. B. Fluctuating and diffusion-limited hypoxia in hypoxia-induced metastasis. *Clin Cancer Res*, 13: 1971-1978, 2007.
262. Shin, K. H., Diaz-Gonzalez, J. A., Russell, J., Chen, Q., Burgman, P., Li, X. F., and Ling, C. C. Detecting Changes in Tumor Hypoxia with Carbonic Anhydrase IX and Pimonidazole. *Cancer Biol Ther*, 6: 70-75, 2007.
263. Silva, P., Homer, J. J., Slevin, N. J., Musgrove, B. T., Sloan, P., Price, P., and West, C. M. Clinical and biological factors affecting response to radiotherapy in patients with head and neck cancer: a review. *Clin Otolaryngol*, 32: 337-345, 2007.
264. Terada, N., Ohno, N., Saitoh, S., and Ohno, S. Immunohistochemical detection of hypoxia in mouse liver tissues treated with pimonidazole using "in vivo cryotechnique". *Histochem Cell Biol*, 128: 252-261, 2007.
265. Thored, P., Wood, J., Arvidsson, A., Cammenga, J., Kokaia, Z., and Lindvall, O. Long-Term Neuroblast Migration Along Blood Vessels in an Area With Transient Angiogenesis and Increased Vascularization After Stroke. *Stroke* 38: 3032-3039, 2007.

266. Tugues, S., Fernandez-Varo, G., Munoz-Luque, J., Ros, J., Arroyo, V., Rodes, J., Friedman, S. L., Carmeliet, P., Jimenez, W., and Morales-Ruiz, M. Antiangiogenic treatment with sunitinib ameliorates inflammatory infiltrate, fibrosis, and portal pressure in cirrhotic rats. *Hepatology*, *46*: 1919-1926, 2007.
267. van der Bilt, J. D., Soeters, M. E., Duyverman, A. M., Nijkamp, M. W., Witteveen, P. O., van Diest, P. J., Kranenburg, O., and Borel Rinkes, I. H. Perinecrotic hypoxia contributes to ischemia/reperfusion-accelerated outgrowth of colorectal micrometastases. *Am J Pathol*, *170*: 1379-1388, 2007.
268. Westbury, C. B., Pearson, A., Nerurkar, A., Reis-Filho, J. S., Steele, D., Peckitt, C., Sharp, G., and Yarnold, J. R. Hypoxia can be detected in irradiated normal human tissue: a study using the hypoxic marker pimonidazole hydrochloride. *Br J Radiol*, *80*: 934-938, 2007.
269. Wijffels, K. I., Marres, H. A., Peters, J. P., Rijken, P. F., van der Kogel, A. J., and Kaanders, J. H. Tumour cell proliferation under hypoxic conditions in human head and neck squamous cell carcinomas. *Oral Oncol*, *44*: 335-344, 2008.
270. Zips, D., Yaromina, A., Schutze, C., Wullrich, K., Krause, M., Krause, M., Hessel, F., Eicheler, W., Dorfler, A., Bruchner, K., Menegakis, A., Zhou, X., Bergmann, R., van den Hoff, J., Beuthien-Baumann, B., and Baumann, M. Experimental evaluation of functional imaging for radiotherapy. *Strahlenther Onkol*, *183 Spec No 2*: 41-42, 2007.
271. Bache, M., Kappler, M., Said, H. M., Staab, A., and Vordermark, D. Detection and specific targeting of hypoxic regions within solid tumors: current preclinical and clinical strategies. *Curr Med Chem*, *15*: 322-338, 2008.
272. Bayer, C., Schilling, D., Hoetzel, J., Egermann, H. P., Zips, D., Yaromina, A., Geurts-Moespot, A., Sprague, L. D., Sweep, F., Baumann, M., Molls, M., and Adam, M. PAI-1 levels predict response to fractionated irradiation in 10 human squamous cell carcinoma lines of the head and neck. *Radiother Oncol*, *86*: 361-368, 2008.
273. Bouchentouf, M., Benabdallah, B. F., Bigey, P., Yau, T. M., Scherman, D., and Tremblay, J. P. Vascular endothelial growth factor reduced hypoxia-induced death of human myoblasts and improved their engraftment in mouse muscles. *Gene Ther*, *15*: 404-414, 2008.
274. Boutrid, H., Jockovich, M. E., Murray, T. G., Pina, Y., Feuer, W. J., Lampidis, T. J., and Cebulla, C. M. Targeting hypoxia, a novel treatment for advanced retinoblastoma. *Invest Ophthalmol Vis Sci*, *49*: 2799-2805, 2008.
275. Busk, M., Horsman, M. R., Jakobsen, S., Keiding, S., van der Kogel, A. J., Bussink, J., and Overgaard, J. Imaging Hypoxia in Xenografted and Murine Tumors With (18)F-Fluoroazomycin Arabinoside: A Comparative Study Involving microPET, Autoradiography, Po(2)-Polarography, and Fluorescence Microscopy. *Int J Radiat Oncol Biol Phys*, *70*: 1202-1212, 2008.
276. Busk, M., Horsman, M. R., Kristjansen, P. E., van der Kogel, A. J., Bussink, J., and Overgaard, J. Aerobic glycolysis in cancers: implications for the usability of oxygen-responsive genes and fluorodeoxyglucose-PET as markers of tissue hypoxia. *Int J Cancer*, *122*: 2726-2734, 2008.
277. Bussink, J., van der Kogel, A. J., and Kaanders, J. H. Patterns and levels of hypoxia in head and neck squamous cell carcinomas and their relationship to

- patient outcome: in regard to Evans et al. (*Int J Radiat Oncol Biol Phys* 2007;69:1024-1031). *Int J Radiat Oncol Biol Phys*, 70: 1616, 2008.
278. Bussink, J., van der Kogel, A. J., and Kaanders, J. H. Activation of the PI3-K/AKT pathway and implications for radioresistance mechanisms in head and neck cancer. *Lancet Oncol*, 9: 288-296, 2008.
  279. Efrati, S., Berman, S., Ben Aharon, G., Siman-Tov, Y., Averbukh, Z., and Weissgarten, J. Application of normobaric hyperoxia therapy for amelioration of haemorrhagic shock-induced acute renal failure. *Nephrol Dial Transplant*, 23: 2213-2222, 2008.
  280. Fujita, N., Imai, J., Suzuki, T., Yamada, M., Ninomiya, K., Miyamoto, K., Iwasaki, R., Morioka, H., Matsumoto, M., Chiba, K., Watanabe, S., Suda, T., Toyama, Y., and Miyamoto, T. Vascular endothelial growth factor-A is a survival factor for nucleus pulposus cells in the intervertebral disc. *Biochem Biophys Res Commun*, 372: 367-372, 2008.
  281. Greijer, A. E., Delis-van Diemen, P. M., Fijneman, R. J., Giles, R. H., Voest, E. E., van Hinsbergh, V. W., and Meijer, G. A. Presence of HIF-1 and related genes in normal mucosa, adenomas and carcinomas of the colorectum. *Virchows Arch*, 452(5):535-544, 2008.
  286. Heyman, S. N., Khamaisi, M., Rosen, S., and Rosenberger, C. Renal Parenchymal Hypoxia, Hypoxia Response and the Progression of Chronic Kidney Disease. *Am J Nephrol*, 28: 998-1006, 2008.
  287. Holcombe, D. J., Lengefeld, N., Gole, G. A., and Barnett, N. L. The effects of acute intraocular pressure elevation on rat retinal glutamate transport. *Acta Ophthalmol*, 86: 408-414, 2008.
  288. Klinkenberg, L. G., Sutherland, L. A., Bishai, W. R., and Karakousis, P. C. Metronidazole Lacks Activity against Mycobacterium tuberculosis in an In Vivo Hypoxic Granuloma Model of Latency. *J Infect Dis*, 198: 275-283, 2008.
  289. Koch, C. J. Importance of antibody concentration in the assessment of cellular hypoxia by flow cytometry: EF5 and pimonidazole. *Radiat Res*, 169: 677-688, 2008.
  290. Kumar, S., Brown, S. L., Kolozsvary, A., Freytag, S. O., and Kim, J. H. Efficacy of suicide gene therapy in hypoxic rat 9L glioma cells. *J Neurooncol*, 2008.
  291. Li, X.-F. and JA, O. D. Hypoxia in microscopic tumors. *Cancer Letters*, 264: 172-180, 2008.
  292. Mayr, M., Sidibe, A., and Zampetaki, A. The paradox of hypoxic and oxidative stress in atherosclerosis. *J Am Coll Cardiol*, 51: 1266-1267, 2008.
  293. Miyazaki, Y., Hara, A., Kato, K., Oyama, T., Yamada, Y., Mori, H., and Shibata, T. The effect of hypoxic microenvironment on matrix metalloproteinase expression in xenografts of human oral squamous cell carcinoma. *Int J Oncol*, 32: 145-151, 2008.
  294. Oh, C., Dong, Y., Harman, C., Mighty, H. E., Kopelman, J., and Thompson, L. P. Chronic hypoxia differentially increases glutathione content and gamma-glutamyl cysteine synthetase expression in fetal guinea pig organs. *Early Hum Dev*, 84: 121-127, 2008.

295. Rausch, M. E., Weisberg, S., Vardhana, P., and Tortoriello, D. V. Obesity in C57BL/6J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration. *Int J Obes (Lond)*, 32: 451-463, 2008.
296. Ream, M. A., Ray, A. M., Chandra, R., and Chikaraishi, D. M. Early Fetal Hypoxia Leads to Growth Restriction and Myocardial Thinning. *Am J Physiol Regul Integr Comp Physiol*, 2008.
297. Rosenberger, C., Goldfarb, M., Shina, A., Bachmann, S., Frei, U., Eckardt, K. U., Schrader, T., Rosen, S., and Heyman, S. N. Evidence for sustained renal hypoxia and transient hypoxia adaptation in experimental rhabdomyolysis-induced acute kidney injury. *Nephrol Dial Transplant*, 23: 1135-1143, 2008.
298. Rosenberger, C., Khamaisi, M., Abassi, Z., Shilo, V., Weksler-Zangen, S., Goldfarb, M., Shina, A., Zibertrest, F., Eckardt, K. U., Rosen, S., and Heyman, S. N. Adaptation to hypoxia in the diabetic rat kidney. *Kidney Int*, 73: 34-42, 2008.
299. Rudolfsson, S. H. and Bergh, A. Testosterone-stimulated growth of the rat prostate may be driven by tissue hypoxia and hypoxia-inducible factor-1alpha. *J Endocrinol*, 196: 11-19, 2008.
300. Saito, Y., Uppal, A., Byfield, G., Budd, S., and Hartnett, M. E. Activated NAD(P)H Oxidase from Supplemental Oxygen Induces Neovascularization Independent of VEGF in Retinopathy of Prematurity Model. *Invest Ophthalmol Vis Sci*, 49: 1591-1598, 2008.
301. Scigliano, S., Pinel, S., Poussier, S., Fouyssac, F., Plenat, F., Karcher, G., and Chastagner, P. Measurement of hypoxia using invasive oxygen-sensitive electrode, pimonidazole binding and 18F-FDG uptake in anaemic or erythropoietin-treated mice bearing human glioma xenografts. *Int J Oncol*, 32: 69-77, 2008.
302. Selvakumaran, M., Yao, K. S., Feldman, M. D., and O'Dwyer, P. J. Antitumor effect of the angiogenesis inhibitor bevacizumab is dependent on susceptibility of tumors to hypoxia-induced apoptosis. *Biochem Pharmacol*, 75: 627-638, 2008.
303. Sersa, G., Jarm, T., Kotnik, T., Coer, A., Podkrajsek, M., Sentjurc, M., Miklavcic, D., Kadivec, M., Kranjc, S., Secerov, A., and Cemazar, M. Vascular disrupting action of electroporation and electrochemotherapy with bleomycin in murine sarcoma. *Br J Cancer*, 98: 388-398, 2008.
304. Sluimer, J. C., Gasc, J. M., van Wanroij, J. L., Kisters, N., Groeneweg, M., Sollewijn Gelpke, M. D., Cleutjens, J. P., van den Akker, L. H., Corvol, P., Wouters, B. G., Daemen, M. J., and Bijnens, A. P. Hypoxia, hypoxia-inducible transcription factor, and macrophages in human atherosclerotic plaques are correlated with intraplaque angiogenesis. *J Am Coll Cardiol*, 51: 1258-1265, 2008.
305. Sun, X., Li, X. F., Russell, J., Xing, L., Urano, M., Li, G. C., Humm, J. L., and Ling, C. C. Changes in tumor hypoxia induced by mild temperature hyperthermia as assessed by dual-tracer immunohistochemistry. *Radiother Oncol*, 2008.
306. Takahashi, M., Yasui, H., Ogura, A., Asanuma, T., Kubota, N., Tsujitani, M., Kuwabara, M., and Inanami, O. X Irradiation Combined with TNF alpha-related Apoptosis-inducing Ligand (TRAIL) Reduces Hypoxic Regions of Human Gastric Adenocarcinoma Xenografts in SCID Mice. *J Radiat Res (Tokyo)*, 49: 153-161, 2008.

307. Troost, E. G., Laverman, P., Philippens, M. E., Lok, J., van der Kogel, A. J., Oyen, W. J., Boerman, O. C., Kaanders, J. H., and Bussink, J. Correlation of [(18)F]FMISO autoradiography and pimonodazole immunohistochemistry in human head and neck carcinoma xenografts. *Eur J Nucl Med Mol Imaging*, 2008.
308. Via, L. E., Lin, P. L., Ray, S. M., Carrillo, J., Allen, S. S., Eum, S. Y., Taylor, K., Klein, E., Manjunatha, U., Gonzales, J., Lee, E. G., Park, S. K., Raleigh, J. A., Cho, S. N., McMurray, D. N., Flynn, J. L., and Barry, C. E., 3rd Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and nonhuman primates. *Infect Immun*, 76: 2333-2340, 2008.
309. Vignozzi, L., Morelli, A., Filippi, S., Vannelli, G. B., Mungai, S., Marini, M., Boddi, V., Forti, G., and Maggi, M. Effect of sildenafil administration on penile hypoxia induced by cavernous neurotomy in the rat. *Int J Impot Res*, 20: 60-67, 2008.